

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00	A2	(11) International Publication Number: WO 00/54759 (43) International Publication Date: 21 September 2000 (21.09.00)
(21) International Application Number: PCT/US00/06611 (22) International Filing Date: 15 March 2000 (15.03.00) (30) Priority Data: 60/124,525 15 March 1999 (15.03.99) US (71) Applicant: TULARIK INC. [US/US]; Two Corporate Drive, South San Francisco, CA 94080 (US). (72) Inventors: LI, Leping; 2100 Davis Drive, Burlingame, CA 94010 (US). MEDINA, Julio, C.; 1407 Cedar St., San Carlos, CA 94070 (US). HASEGAWA, Hirohiko; 1-98, 1 Kasugadenaka 3-chome, Konohana-ku, Osaka 554 (JP). CUTLER, Serena, T.; 449 Hawthorne Ave. #3, Palo Alto, CA 94301 (US). LIU, Jiwen; Apt. 211, 1075 Rollins Rd., Burlingame, CA 94010 (US). ZHU, Liusheng; Apt. 211, 2001 Trousdale Drive, Burlingame, CA 94010 (US). SHAN, Bei; 349 Quay Lane, Redwood City, CA 94065 (US). LUSTIG, Kevin; Two Corporate Drive, South San Francisco, CA 94080 (US). (74) Agents: KEZER, William, B. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: LXR MODULATORS (57) Abstract The invention provides compounds, compositions and methods for modulating the effects of LXR in a cell. The compounds and compositions are useful both as diagnostic indicators of LXR function and as pharmacologically active agents. The compounds and compositions find particular use in the treatment of disease states associated with cholesterol metabolism, particularly atherosclerosis and hypercholesterolemia.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

LXR MODULATORS

FIELD OF THE INVENTION

5

The present invention relates to compounds and methods useful for the modulation of LXR. In view of the activity of LXR in the control of cholesterol homeostasis, the compounds described herein are useful for lowering plasma cholesterol levels.

10

BACKGROUND

Cholesterol is used for the synthesis of bile acids in the liver, the manufacture and repair of cell membranes, and the synthesis of steroid hormones.

15 There are both exogenous and endogenous sources of cholesterol. The average American consumes about 450 mg of cholesterol each day and produces an additional 500 to 1,000 mg in the liver and other tissues. Another source is the 500 to 1,000 mg of biliary cholesterol that is secreted into the intestine daily; about 50 percent is reabsorbed (enterohepatic circulation). Excess accumulation of cholesterol in the
20 arterial walls can result in atherosclerosis, which is characterized by plaque formation. The plaques inhibit blood flow, promote clot formation and can ultimately cause heart attacks, stroke and claudication. Development of therapeutic agents for the treatment of atherosclerosis and other diseases associated with cholesterol metabolism has been focused on achieving a more complete understanding of the biochemical pathways
25 involved. Most recently, liver X receptors (LXRs) were identified as key components in cholesterol homeostasis.

The LXRs were first identified as orphan members of the nuclear receptor superfamily whose ligands and functions were unknown. Two LXR proteins (α and β) are known to exist in mammals. The expression of LXR α is restricted, with
30 the highest levels being found in the liver, and lower levels found in kidney, intestine, spleen, and adrenals (see Willy, *et al.*, *Genes Dev.* **9**(9):1033-45 (1995)). LXR β is rather ubiquitous, being found in nearly all tissues examined. Recent studies on the LXRs indicate that they are activated by certain naturally occurring, oxidized derivatives of cholesterol, including 22(R)-hydroxycholesterol, 24(S)-
35 hydroxycholesterol and 24,25(S)-epoxycholesterol (see Lehmann, *et al.*, *J. Biol. Chem.* **272**(6):3137-3140 (1997)). The expression pattern of LXRs and their

oxysterol ligands provided the first hint that these receptors may play a role in cholesterol metabolism (see Janowski, *et al.*, *Nature* 383:728-731 (1996)).

As noted above, cholesterol metabolism in mammals occurs via conversion into steroid hormones or bile acids. The role of LXRs in cholesterol homeostasis was first postulated to involve the pathway of bile acid synthesis, in which cholesterol 7 α -hydroxylase (CYP7A) operates in a rate-limiting manner. Support for this proposal was provided when additional experiments found that the CYP7A promoter contained a functional LXR response element that could be activated by RXR/LXR heterodimers in an oxysterol- and retinoid-dependent manner. Confirmation of LXR function as a transcriptional control point in cholesterol metabolism was made using knockout mice, particularly those lacking the oxysterol receptor LXR α (see Peet, *et al.*, *Cell* 93:693-704 (1998)).

Mice lacking the receptor LXR α (*e.g.*, knockout or (-/-) mice) lost their ability to respond normally to increases in dietary cholesterol and were unable to tolerate any cholesterol in excess of that synthesized *de novo*. LXR α (-/-) mice did not induce transcription of the gene encoding CYP7A when fed diets containing additional cholesterol. This resulted in an accumulation of large amounts of cholesterol and impaired hepatic function in the livers of LXR α (-/-) mice. These results further established the role of LXR α as the essential regulatory component of cholesterol homeostasis. LXR α is also believed to be involved in fatty acid synthesis. Accordingly, regulation of LXR α (*e.g.*, use of LXR α agonist or antagonists) could provide treatment for a variety of lipid disorders including obesity and diabetes.


In view of the importance of LXRs, and particularly LXR α s to the delicate balance of cholesterol metabolism and fatty acid biosynthesis, we describe modulators of LXRs which are useful as therapeutic agents or diagnostic agents for the treatment of disorders associated with bile acid and cholesterol metabolism, including cholesterol gallstones, atherosclerosis, lipid storage diseases, obesity, and diabetes. The agents described herein are also useful for disease states associated with serum hypercholesterolemia, such as coronary heart disease.

30

SUMMARY OF THE INVENTION

In one aspect, the present invention provides compositions for modulation of LXR α function in a cell. The compositions typically comprise a pharmaceutically acceptable excipient and a compound having the formula:

35



(I)

wherein Ar represents an aryl group; R¹ is -OH, -O-(C₁-C₇)alkyl, -OC(O)-(C₁-C₇)alkyl, -O-(C₁-C₇)heteroalkyl, -OC(O)-(C₁-C₇)heteroalkyl, -CO₂H, -NH₂, -NH(C₁-C₇)alkyl, -N((C₁-C₇)alkyl)₂ or -NH-S(O)₂-(C₁-C₃)alkyl; R² is (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl; X¹, X², X³, X⁴, X⁵ and X⁶ are each independently H, (C₁-C₃)alkyl, (C₁-C₃)heteroalkyl, F or Cl, with the proviso that no more than three of X¹ through X⁶ are H, (C₁-C₃)alkyl or (C₁-C₃)heteroalkyl; and Y is -N(R¹²)S(O)_m-, -N(R¹²)S(O)_mN(R¹³)-, -N(R¹²)C(O)-, -N(R¹²)C(O)N(R¹³)-, -N(R¹²)C(S)- or -N(R¹²)C(O)O-, wherein R¹² and R¹³ are each independently hydrogen, (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl, and optionally when Y is -N(R¹²)S(O)_m- or -N(R¹²)S(O)_mN(R¹³)-, R¹² forms a five-, six- or seven-membered ring fused to Ar or to R² through covalent attachment to Ar or R², respectively. In the above Y groups, the subscript m is an integer of from 1 to 2.

Preferred compositions are those in which the compound above binds to the ligand binding domain of LXRα with an affinity of at least 1 micromolar.

A number of compounds that are useful in the above-described compositions are novel. Accordingly, the present invention further provides compounds of the above formula, with the proviso that when R¹ is OH, and -Y-R² is -N(R¹²)S(O)_m-R² or -N(R¹²)C(O)N(R¹³)-R² and is attached to a position para to the quaternary carbon attached to Ar, and when R² is phenyl, benzyl or benzoyl, then i) at least one of R¹² or R¹³ is other than hydrogen and contains an electron-withdrawing substituent, or ii) R² is substituted with a moiety other than amino, acetamido, di(C₁-C₇)alkylamino, (C₁-C₇)alkylamino, halogen, hydroxy, nitro, or (C₁-C₇)alkyl, or iii) the benzene ring portion of R² is trisubstituted in addition to the Y group.

In yet another aspect, the present invention provides methods for modulating LXR in a cell by administering to or contacting the cell with a composition containing a compound of Formula I above.

In still another aspect, the present invention provides methods for treating LXR-responsive diseases by administering to a subject in need of such treatment a composition containing a compound of Formula I. These methods are particularly useful for the treatment of pathology such as hypercholesterolemia, atherosclerosis, and hyperlipoproteinemia. In certain embodiments, the compound can be administered to the subject in combination with an additional anti-

hypercholesterolemic agent, for example, bile acid sequestrants, nicotinic acid, fibric acid derivatives or HMG CoA reductase inhibitors.

The present compounds can exert their effects either systemically (the compounds permeate the relevant tissues, such as liver, upon entrance into the bloodstream) or locally (for example, by modulating LXR function of intestinal epithelial cells following oral administration, without necessitating the compounds' entrance into the bloodstream). In some disease states, some preferred compounds will be those with good systemic distribution, while, in other instances, preferred compounds will be those that can work locally on the intestinal track or on the skin without penetrating the bloodstream.

Certain compounds of the present invention are antiproliferative and can be used in compositions for treating diseases associated with abnormal cell proliferation (*e.g.*, cancer). Other diseases associated with an abnormally high level of cellular proliferation include restenosis, where vascular smooth muscle cells are involved, inflammatory disease states, where endothelial cells, inflammatory cells and glomerular cells are involved, myocardial infarction, where heart muscle cells are involved, glomerular nephritis, where kidney cells are involved, transplant rejection, where endothelial cells are involved, infectious diseases such as HIV infection and malaria, where certain immune cells and/or other infected cells are involved, and the like. Infectious and parasitic agents *per se* (*e.g.* bacteria, trypanosomes, fungi, *etc*) are also subject to selective proliferative control using the subject compositions and compounds.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1 and 2 provide the structures of compounds prepared using the solid phase synthesis methodology outlined in Scheme 3.

30

DETAILED DESCRIPTION OF THE INVENTION

Definitions

The term "alkyl," by itself or as part of another substituent, means, unless otherwise stated, a straight or branched chain, or cyclic hydrocarbon radical, or combination thereof, which may be fully saturated, mono- or polyunsaturated and can

include di- and multi-radicals, having the number of carbon atoms designated (*i.e.* C₁-C₁₀ means one to ten carbons). Examples of saturated hydrocarbon radicals include groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, cyclohexyl, (cyclohexyl)methyl, cyclopropylmethyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butynyl, and the higher homologs and isomers. The term "alkyl," unless otherwise noted, is also meant to include those derivatives of alkyl defined in more detail below as "cycloalkyl" and "alkylene." The term "alkylene" by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified by -CH₂CH₂CH₂CH₂-. Typically, an alkyl group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the present invention. A "lower alkyl" or "lower alkylene" is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms, preferably four or fewer carbon atoms.

The term "alkoxy," employed alone or in combination with other terms means, unless otherwise stated, an alkyl group, as defined above, connected to the remainder of the molecule via an oxygen atom, such as, for example, methoxy, ethoxy, 1-propoxy, 2-propoxy and the higher homologs and isomers.

The term "heteroalkyl," by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, consisting of the stated number of carbon atoms and from one to three heteroatoms selected from the group consisting of O, N, Si and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) O, N and S may be placed at any position of the heteroalkyl group except for the position at which the heteroalkyl group is attached to the rest of the molecule. The heteroatom Si may be placed at any position of the heteroalkyl group, including the position at which the alkyl group is attached to the remainder of the molecule. Examples include -CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, and -CH=CH-N(CH₃)-CH₃. Up to two heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. Also included in the term "heteroalkyl" are those radicals described in more detail below as "heteroalkylene" and "heterocycloalkyl." The term "heteroalkylene" by itself or as part of another

substituent means a divalent radical derived from heteroalkyl, as exemplified by $-\text{CH}_2-\text{CH}_2-\text{S}-\text{CH}_2\text{CH}_2-$ and $-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_2-$. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini. Still further, for alkylene and heteroalkylene linking groups, as well as all other linking groups
5 described herein, no specific orientation of the linking group is implied.

The terms "cycloalkyl" and "heterocycloalkyl", by themselves or in combination with other terms, represent, unless otherwise stated, cyclic versions of "alkyl" and "heteroalkyl", respectively. The terms "cycloalkyl" and "heterocycloalkyl" are also meant to include bicyclic, tricyclic and polycyclic versions
10 thereof. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl, adamantyl, and the like. Examples of heterocycloalkyl include 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-
15 piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, 1,4-diazabicyclo[2.2.2]oct-2-yl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like.

The terms "halo" or "halogen," by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine
20 atom. Additionally, terms such as "fluoroalkyl," are meant to include monofluoroalkyl and polyfluoroalkyl.

The term "aryl," employed alone or in combination with other terms (e.g., aryloxy, arylthioxy, arylalkyl) means, unless otherwise stated, an aromatic substituent which can be a single ring or multiple rings (up to three rings) which are
25 fused together or linked covalently. The rings may each contain from zero to four heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. The aryl groups that contain heteroatoms may be referred to as "heteroaryl" and can be attached to the remainder of the molecule through a carbon atom or a heteroatom.
30 Non-limiting examples of aryl groups include phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl,
35 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxaliny, 5-quinoxaliny, 3-quinolyl, and 6-quinolyl. Substituents for each of the above noted

aryl ring systems are selected from the group of acceptable substituents described below.

The terms "arylalkyl" and "arylheteroalkyl" are meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, phenethyl, pyridylmethyl and the like) or a heteroalkyl group (*e.g.*, phenoxymethyl, 2-pyridyloxymethyl, 1-naphthyloxy-3-propyl, and the like). The arylalkyl and arylheteroalkyl groups will typically contain from 1 to 3 aryl moieties attached to the alkyl or heteroalkyl portion by a covalent bond or by fusing the ring to, for example, a cycloalkyl or heterocycloalkyl group. For arylheteroalkyl groups, a heteroatom can occupy the position at which the group is attached to the remainder of the molecule. For example, the term "arylheteroalkyl" is meant to include benzyloxy, 2-phenylethoxy, phenethylamine, and the like.

Each of the above terms (*e.g.*, "alkyl," "heteroalkyl," "aryl," *etc.*) are meant to include both substituted and unsubstituted forms of the indicated radical.

Preferred substituents for each type of radical are provided below.

Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be a variety of groups selected from: -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)NR'R''', -NR''C(O)₂R', -NHC(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -CN and -NO₂ in a number ranging from zero to (2N+1), where N is the total number of carbon atoms in such radical. Preferably, substituted alkyl groups will have from one to six independently selected substituents, more preferably from one to four independently selected substituents, most preferably from one to three independently selected substituents. In the substituents listed above, R', R'' and R''' each independently refer to hydrogen, unsubstituted(C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl, aryl substituted with 1-3 halogens, unsubstituted alkyl, alkoxy or thioalkoxy groups, or aryl-(C₁-C₄)alkyl groups. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-membered ring. For example, -NR'R'' is meant to include 1-pyrrolidinyl and 4-morpholinyl.

Similarly, substituents for the aryl groups are varied and are selected from: -halogen, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR''C(O)NR'R''', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -N₃, -CH(Ph)₂, perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in a number ranging

from zero to the total number of open valences on the aromatic ring system; and where R' and R'' are independently selected from hydrogen, (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl, (unsubstituted aryl)-(C₁-C₄)alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl. Preferably, substituted aryl groups will have from one to four
5 independently selected substituents, more preferably from one to three independently selected substituents, most preferably from one to two independently selected substituents.

Two of the substituents on adjacent atoms of the aryl ring may optionally be replaced with a substituent of the formula -T-C(O)-(CH₂)_q-U-, wherein T
10 and U are independently -NH-, -O-, -CH₂- or a single bond, and q is an integer of from 0 to 2. Alternatively, two of the substituents on adjacent atoms of the aryl ring may optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein A and B are independently -CH₂-, -O-, -NH-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single bond, and r is an integer of from 1 to 3. One of the single bonds of the new ring so
15 formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl ring may optionally be replaced with a substituent of the formula -(CH₂)_s-X-(CH₂)_t-, where s and t are independently integers of from 0 to 3, and X is -O-, -NR'-, -S-, -S(O)-, -S(O)₂-, or -S(O)₂NR'-. The substituent R' in -NR'- and -S(O)₂NR'- is selected from hydrogen or unsubstituted
20 (C₁-C₆)alkyl.

As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

The term "pharmaceutically acceptable salts" is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases,
25 depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium,
30 potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived
35 from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric,

dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, oxalic, maleic, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (see, for example, Berge, S.M., et al, "Pharmaceutical Salts", *Journal of Pharmaceutical Science*, 1977, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

In addition to salt forms, the present invention provides compounds which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide a compound of formula I. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the present invention when placed in a transdermal patch reservoir with a suitable enzyme.

Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers and individual isomers are all intended to be encompassed within the scope of the present invention.

The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such

compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be encompassed within the scope of the present invention.

5

General


The present invention provides compositions, compounds and methods for modulating LXR function in a cell. The compositions which are useful for this modulation will typically be those which contain an effective amount of an LXR-modulating compound. In general, an effective amount of an LXR-modulating compound is a concentration of the compound that will produce at 50 percent increase/decrease in LXR activity in a cell-based reporter gene assay, or a biochemical peptide-sensor assay such as the assays described in co-pending applications Ser. Nos. 08/975,614 (filed November 21, 1997) and 09/163,713 (filed September 30, 1998).

15

Embodiments of the Invention

In one aspect, the present invention provides compositions for modulation of LXR function in a cell. The compositions typically comprise a pharmaceutically acceptable excipient and an LXR-modulating amount of a compound having the formula:

20



(I)

The symbol Ar represents an aryl group. A variety of aryl groups, both unsubstituted and substituted, are useful in the present invention. Preferred Ar groups are either monocyclic or fused-bicyclic aromatic rings. Particularly preferred Ar ring systems are benzene, naphthalene, pyridine, quinoline, isoquinoline, pyrrole, furan and thiophene. More preferably, Ar represents either a benzene or pyridine ring, with benzene being the most preferred.

30

When Ar represents a substituted aromatic ring (substituents being in addition to -Y-R² and the carbon bearing R¹), the substituents will typically be selected from -OH, -NH₂, lower alkyl (e.g., methyl, butyl, trifluoromethyl,

trifluoroethyl, and the like), lower alkoxy (e.g., methoxy, ethoxy, trifluoromethoxy, butoxy, and the like), $-NR'R''$, $-SR'$, $-halogen$, $-SiR'R''R'''$, $-OC(O)R'$, $-CO_2R'$, $-CONR'R''$, $-OC(O)NR'R''$, $-NR''C(O)R'$, $-NR''C(O)_2R'$, $-NR''C(O)NR'R'''$, $-NH-C(NH_2)=NH$, $-NR'C(NH_2)=NH$, $-NH-C(NH_2)=NR'$, $-S(O)R'$, $-S(O)_2R'$, $-S(O)_2NR'R''$, $-CN$ and $-NO_2$; wherein R' , R'' and R''' are each independently selected from the group consisting of hydrogen and $(C_1-C_3)alkyl$. Preferred compounds are those in which the number of substituents on the Ar group (in addition to $-Y-R^2$ and the carbon bearing R^1) ranges from 0 to 2.

The groups attached to Ar can be in any spatial arrangement. When Ar is a benzene ring, the two groups shown in Formula I will preferably be attached to Ar in a 1,3-orientation (meta) or a 1,4-orientation (para). More preferably the two groups shown in Formula I will be attached to a benzene or pyridine ring in a 1,4-orientation (para).

Returning now to Formula I, R^1 represents $-OH$, $-O-(C_1-C_7)alkyl$, $-OC(O)-(C_1-C_7)alkyl$, $-O-(C_1-C_7)heteroalkyl$, $-OC(O)-(C_1-C_7)heteroalkyl$, $-CO_2H$, $-NH_2$, $-NH(C_1-C_7)alkyl$, $-N((C_1-C_7)alkyl)_2$ or $-NH-S(O)_2-(C_1-C_5)alkyl$. More preferably, R^1 represents $-OH$, $-CO_2H$, $-NH_2$, $-NH(C_1-C_7)alkyl$, $-N((C_1-C_7)alkyl)_2$ or $-NH-S(O)_2-(C_1-C_5)alkyl$. Most preferably, R^1 is $-OH$. For those embodiments in which R^1 is a dialkylamino group ($-N((C_1-C_7)alkyl)_2$), the alkyl groups can either be the same or different.

The symbols X^1 , X^2 , X^3 , X^4 , X^5 and X^6 each independently represent H, $(C_1-C_5)alkyl$, $(C_1-C_5)heteroalkyl$, F or Cl, with the proviso that no more than three of X^1 through X^6 are H, $(C_1-C_5)alkyl$ or $(C_1-C_5)heteroalkyl$. More preferably, no more than three of X^1 through X^6 are H with the remaining being F. Most preferably, X^1 through X^6 are each F.


The symbol Y represents a linking group selected from $-N(R^{12})S(O)_m-$, $-N(R^{12})S(O)_mN(R^{13})-$, $-N(R^{12})C(O)-$, $-N(R^{12})C(O)N(R^{13})-$, $-N(R^{12})C(S)-$ or $-N(R^{12})C(O)O-$, wherein R^{12} and R^{13} are each independently H, $(C_1-C_7)alkyl$, $(C_1-C_7)heteroalkyl$, aryl and aryl $(C_1-C_7)alkyl$, and optionally when Y is $-N(R^{12})S(O)_m-$ or $-N(R^{12})S(O)_mN(R^{13})-$, R^{12} forms a five-, six- or seven-membered ring fused to Ar or to R^2 through covalent attachment to Ar or to R^2 , respectively. In the above Y groups, the subscript m is an integer of from 1 to 2. Preferably, Y is $-N(R^{12})S(O)_m-$, $-N(R^{12})S(O)_mN(R^{13})-$ or $-N(R^{12})C(O)O-$. Most preferably, Y is $-N(R^{12})S(O)_m-$.

As noted above, R^{12} and R^{13} represent hydrogen, $(C_1-C_7)alkyl$, $(C_1-C_7)heteroalkyl$, aryl or aryl $(C_1-C_7)alkyl$, which in the case of the latter two groups, can also be either substituted or unsubstituted. In one group of embodiments, R^{12} is hydrogen or $(C_1-C_4)alkyl$, preferably fluoro $(C_1-C_4)alkyl$. A particularly preferred R^{12}

group is 2,2,2-trifluoroethyl. In another group of preferred embodiments, R^{12} is attached to Ar to form a fused ring system, for example, indoline, tetrahydroquinoline or tetrahydroisoquinoline.


Attached to Y is R^2 , which represents (C_1-C_7) alkyl, (C_1-C_7) heteroalkyl, aryl or aryl (C_1-C_7) alkyl. Each of the recited R^2 groups can be either substituted or unsubstituted. In preferred embodiments, R^2 is an aryl group. More preferably, R^2 is phenyl, thienyl, imidazolyl, oxazolyl or pyridyl. Most preferably, R^2 is phenyl or thienyl (including 2-thienyl and 3-thienyl). Preferred substituted R^2 groups include 3-chlorophenyl, 3-bromophenyl, 3-cyanophenyl, 3-(trifluoromethyl)phenyl, 2-chloro-3-thienyl and 2,5-dichloro-3-thienyl.

The above recitation describes a number of preferred groups for the compounds and compositions of the present invention. Additionally, certain combinations of the above groups will also be preferred. For example, in one group of embodiments, the compositions of the present invention will include a compound having a formula selected from the group consisting of




In each of the above formulae, the subscript n represents an integer of from 0 to 4; and each R^{11} is independently -OH, $-NH_2$, lower alkyl, lower alkoxy, $-NR'R''$, $-SR'$, -halogen, $-SiR'R''R'''$, $-OC(O)R'$, $-CO_2R'$, $-CONR'R''$, $-OC(O)NR'R''$, $-NR''C(O)R'$, $-NR''C(O)_2R'$, $-NR''C(O)NR'R'''$, $-NH-C(NH_2)=NH$, $-NR'C(NH_2)=NH$, $-NH-C(NH_2)=NR'$, $-S(O)R'$, $-S(O)_2R'$, $-S(O)_2NR'R''$, $-CN$ and $-NO_2$; wherein each R' , R'' and R''' is independently hydrogen or (C_1-C_5) alkyl. The remaining groups in the above formulae have the meanings provided for Formula I.

In a further preferred group of embodiments, the compositions will include a compound having a formula selected from the group consisting of




- In this group of preferred embodiments, the various groups (*e.g.*, R¹, X¹, X⁶, R², R¹¹, R¹² and R¹³) have the meanings indicated above with reference to Formula I. Preferably, R¹ is -OH or -NH₂, and X¹ and X⁶ are each independently hydrogen or fluorine. Still further preferred within this group of embodiments are those in which R² is a substituted aryl group, more preferably a substituted phenyl or substituted thienyl group.

More preferably, the compositions will include a compound having a formula selected from the group consisting of



10

Most preferred, are those embodiments in which the compound has the formula:




Within the group of preferred compounds provided as Formulae IIa or IIb, at least three groupings of substituents are particularly preferred.

In a first group of preferred substituents, R^1 is -OH or -NH₂; X^1 and X^6 are each independently hydrogen or fluorine, R^{12} is fluoro(C₁-C₄)alkyl; and R^2 is aryl.

5 In a second group of preferred substituents, R^1 is -OH or -NH₂; X^1 and X^6 are each independently hydrogen or fluorine, R^{12} is hydrogen or (C₁-C₄)alkyl; and R^2 is substituted or unsubstituted thienyl.

In a third group of preferred substituents, the compositions will include at least one compound of formula II in which R^1 is -OH or -NH₂; X^1 and X^6 are each independently hydrogen or fluorine, R^{12} is (C₁-C₄)alkyl or (C₁-C₄)heteroalkyl; and R^2 is phenyl substituted with at least one member selected from the group consisting of halogen, -CN, -CF₃, -O-(C₁-C₄)alkyl, -C(O)-(C₁-C₄)alkyl, -C(O)-O(C₁-C₄)alkyl, -C(O)-NH(C₁-C₄)alkyl and -C(O)N((C₁-C₄)alkyl)₂. Still further preferred are those
10 embodiments in which R^1 is -OH; and R^2 is phenyl substituted with at least one member selected from -CN, -CF₃ and -O-(C₁-C₄)alkyl.
15

Still other preferred embodiments have a formula selected from:



In each of the formula above, the groups Ar, R^2 , R^{12} , and the subscript m have the meanings provided above with reference to formula I.

20 Further preferred compositions are those in which the compound above binds to the ligand binding domain of an LXR receptor, more preferably LXR α , with an affinity of at least 1 micromolar.

A number of the compounds used in the present compositions are novel. Accordingly, the present invention provides, in another aspect, compounds of
25 Formula I, above, (wherein each of the recited substituents has the meaning provided above) with the proviso that when -Y- R^2 is -N(R^{12})S(O)_m- R^2 or -N(R^{12})C(O)N(R^{13})- R^2 and is attached to a position para to the quaternary carbon attached to Ar, and when R^2 is substituted or unsubstituted phenyl, benzyl or benzoyl, then i) at least one of R^{12} or R^{13} is other than hydrogen or unsubstituted alkyl, or ii) R^2 is substituted with a moiety
30 other than amino, acetamido, di(C₁-C₇)alkylamino, (C₁-C₇)alkylamino, halogen, hydroxy, nitro, or (C₁-C₇)alkyl, or iii) the R^2 phenyl group is trisubstituted in addition to the Y group.

Some of the compounds of Formula I may exist as stereoisomers, and the invention includes all active stereoisomeric forms of these compounds. In the case
35 of optically active isomers, such compounds may be obtained from corresponding

optically active precursors using the procedures described above or by resolving racemic mixtures. The resolution may be carried out using various techniques such as chromatography, repeated recrystallization of derived asymmetric salts, or derivatization, which techniques are well known to those of ordinary skill in the art.

5 The compounds of the invention may be labeled in a variety of ways. For example, the compounds may contain radioactive isotopes such as, for example, ^3H (tritium) and ^{14}C (carbon-14). Similarly, the compounds may be advantageously joined, covalently or noncovalently, directly or through a linker molecule, to a wide variety of other compounds, which may provide pro-drugs or function as carriers,
10 labels, adjuvants, coactivators, stabilizers, *etc.* Such labeled and joined compounds are contemplated within the present invention.


 In yet another aspect, the present invention provides a method for modulating the action of an LXR receptor, preferably LXR α , in a cell. According to this method, the cell is contacted with a sufficient concentration of a composition
15 containing a compound of formula I for either an agonistic or antagonistic effect to be detected. In preferred embodiments, the composition contains an amount of the compound which has been determined to provide a desired therapeutic or prophylactic effect for a given LXR-mediated condition.

 In still another aspect, the present invention provides methods for the
20 treatment of pathology such as hypercholesterolemia, atherosclerosis, and hyperlipoproteinemia using pharmaceutical compositions containing compounds of the foregoing description of the general Formula I. Briefly, this aspect of the invention involves administering to a patient an effective formulation of one or more of the subject compositions. In other embodiments, the compound of Formula I can
25 be administered in combination with other anti-hypercholesterolemic agents (*e.g.*, a bile acid sequestrant, nicotinic acid, fibric acid derivatives or HMG CoA reductase inhibitors), or in combination with other agents that affect cholesterol or lipid metabolism.

30 Synthesis

 Compounds of the present invention can be prepared using readily available starting materials or known intermediates. Scheme 1 provides a variety of synthesis avenues for the production of the subject compounds. One of skill in the art
35 will understand that additional methods are also useful.

SCHEME 1




As shown in Scheme 1, aniline i (as representative of substituted anilines and other arylamines) can either be alkylated, acylated or arylated (general addition of R group) to form ii, or the aromatic ring can be derivatized with, for example, hexafluoroacetone to form iii. Treatment of iii with an appropriate alkylating group, acylating group or arylating group provides iv, which can be sulfonated with, for example, an appropriate sulfonyl halide to form vi. Alternatively, the aniline derivative iii can be sulfonated to form v, which can then be alkylated or acylated to form compounds of formula vi.

Other compounds of the present invention can be formed by treating the substituted aniline iv (or, alternatively, iii), with reagents suitable for the formation of amides vii, carbamates viii, and ureas ix. A variety of reagents are useful in the above scheme and can be found in, for example, March, *Advanced Organic Chemistry* 4th Ed., John Wiley & Sons, New York, NY (1992). Preferred reagents and conditions are also found in the Examples below.

Methods for the preparation of compounds in which a sulfonamide linkage has the reverse orientation are provided in Scheme 2. According to Scheme 2, a substituted benzenesulfonamide **x** (prepared from a substituted benzenesulfonyl chloride and an amine or an aniline) can be converted to the desired alcohol **xi** upon treatment with $\text{CF}_3\text{-TMS}$ in the presence of tetrabutylammonium fluoride in tetrahydrofuran. Alternatively, a halo-substituted benzenesulfonamide **xii** (similarly prepared from a halo-substituted benzenesulfonyl chloride and an amine or an aniline) can be converted to the trifluoromethylketone **xiii** upon treatment with *n*-butyllithium followed by methyl trifluoroacetate at -78°C . Subsequent treatment of **xiii** with $\text{CF}_3\text{-TMS}$ in the presence of tetrabutylammonium fluoride in tetrahydrofuran provides compounds of formula **xi**.

SCHEME 2




Solid Phase Synthesis

Scheme 3 provides one method for preparing compounds of the present invention using solid phase synthesis. In this method, an initial compound is further elaborated using a Suzuki coupling to include additional aryl substituents on a phenyl ring adjacent to the sulfonyl moiety. According to the method outlined in Scheme 2, an activated resin (e.g., 4-(bromomethyl)phenoxyethyl polystyrene from Nova Biochem) is treated with a parent compound to be derivatized (e.g., compound **xiv**) to provide a tethered reactive species **xv**. Suzuki coupling conditions are then employed to attach an aromatic moiety (Ar) in place of the bromine substituent (see Step 2) and provide a modified target compound tethered to the support (**xvi**). The modified target compound (**xvii**) is then cleaved from the support and filtered through silica gel.

Figures 1 and 2 provide structures of compounds prepared using this methodology. The aryl boronic acids used in the Suzuki coupling were from commercial sources (e.g., Aldrich Chemical Co., Milwaukee, Wisconsin, USA; Lancaster, Digital Specialty Chemicals Inc., and Combi-Blocks).

5

SCHEME 3



10

Analysis of compounds

Representative compounds and compositions were demonstrated to have pharmacological activity in *in vitro* and *in vivo* assays, e.g., they are capable of specifically modulating a cellular physiology to reduce an associated pathology or provide or enhance a prophylaxis.

Certain preferred compounds and compositions are capable of specifically regulating LXR. Compounds may be evaluated *in vitro* for their ability to activate LXR receptor function using biochemical assays (see co-pending applications Ser. Nos. 08/975,614 (filed November 21, 1997) and 09/163,713 (filed September 30,

1998)), or in cell-based assays such as that described in Lehnman, et al. (*J. Biol. Chem.* **1997**, 272(6), 3137-3140). Alternatively, the compounds and compositions can be evaluated for their ability to increase or decrease gene expression modulated by LXR, using western-blot analysis. Established animal models to evaluate
5 hypocholesterolemic effects of the compounds are also known in the art. For example, compounds disclosed herein can lower cholesterol levels in hamsters fed a high-cholesterol diet, using a protocol similar to that described in Spady et al. (*J. Clin. Invest.* **1988**, 81, 300), Evans et al. (*J. Lipid Res.* **1994**, 35, 1634), and Lin et al. (*J. Med. Chem.* **1995**, 38, 277). Still further, LXR α animal models (*e.g.*, LXR α (+/-) and (-/-) mice) can be used for evaluation of the present compounds and compositions
10 (see, for example, Peet, et al. *Cell* **1998**, 93, 693-704).

Accordingly, as used herein, the term "LXR-modulating amount" refers to that amount of a compound that is needed to produce a desired effect in any one of the cell-based assays, biochemical assays or animal models described above.
15 Typically, an LXR-modulating amount of a compound will be at least that amount which exhibits an EC₅₀ in a reporter-gene cell-based assay (relative to an untreated control).

Formulation and administration of compounds and pharmaceutical compositions

20

The invention provides methods of using the subject compounds and compositions to treat disease or provide medicinal prophylaxis, to activate LXR receptor function in a cell, to reduce blood cholesterol concentration in a host, to slow down and/or reduce the abnormal cellular proliferation including the growth of
25 tumors, *etc.* These methods generally involve contacting the cell or cells with or administering to a host an effective amount of the subject compounds or pharmaceutically acceptable compositions.

The compositions and compounds of the invention and the pharmaceutically acceptable salts thereof can be administered in any effective way
30 such as via oral, parenteral or topical routes. Generally, the compounds are administered in dosages ranging from about 2 mg up to about 2,000 mg per day, although variations will necessarily occur depending on the disease target, the patient, and the route of administration. Preferred dosages are administered orally in the range of about 0.05 mg/kg to about 20 mg/kg, more preferably in the range of about 0.05
35 mg/kg to about 2 mg/kg, most preferably in the range of about 0.05 mg/kg to about 0.2 mg per kg of body weight per day.

In one embodiment, the invention provides the subject compounds combined with a pharmaceutically acceptable excipient such as sterile saline or other medium, water, gelatin, an oil, *etc.* to form pharmaceutically acceptable compositions. The compositions and/or compounds may be administered alone or in combination
5 with any convenient carrier, diluent, *etc.* and such administration may be provided in single or multiple dosages. Useful carriers include solid, semi-solid or liquid media including water and non-toxic organic solvents.

In another embodiment, the invention provides the subject compounds in the form of a pro-drug, which can be metabolically converted to the subject
10 compound by the recipient host. A wide variety of pro-drug formulations are known in the art.

The compositions may be provided in any convenient form including tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, suppositories, *etc.* As such the compositions, in pharmaceutically acceptable dosage
15 units or in bulk, may be incorporated into a wide variety of containers. For example, dosage units may be included in a variety of containers including capsules, pills, *etc.*

The compositions may be advantageously combined and/or used in combination with other hypocholesterolemic therapeutic or prophylactic agents, different from the subject compounds. In many instances, administration in
20 conjunction with the subject compositions enhances the efficacy of such agents. Exemplary hypocholesterolemic and/or hypolipemic agents include: bile acid sequestrants such as quaternary amines (*e.g.* cholestyramine and colestipol); nicotinic acid and its derivatives; HMG-CoA reductase inhibitors such as mevastatin, pravastatin, and simvastatin; gemfibrozil and other fibric acids, such as clofibrate,
25 fenofibrate, benzaifibrate and cipofibrate; probucol; raloxifene and its derivatives; and mixtures thereof.

The compounds and compositions also find use in a variety of *in vitro* and *in vivo* assays, including diagnostic assays. For example, various allotypic LDL receptor gene expression processes may be distinguished in sensitivity assays with the
30 subject compounds and compositions, or panels thereof. In certain assays and in *in vivo* distribution studies, it is desirable to use labeled versions of the subject compounds and compositions, *e.g.* radioligand displacement assays. Accordingly, the invention provides the subject compounds and compositions comprising a detectable label, which may be spectroscopic (*e.g.* fluorescent), radioactive, *etc.*

35


The following examples are offered by way of illustration and not by way of limitation.

EXAMPLES

¹H-NMR spectra were recorded on a Varian Gemini 400 MHz NMR spectrometer. Significant peaks are tabulated and typically include: number of protons, multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br s, broad singlet) and coupling constant(s) in Hertz. Electron Ionization (EI) mass spectra were recorded on a Hewlett Packard 5989A mass spectrometer. Mass spectrometry results are reported as the ratio of mass over charge, followed by the relative abundance of each ion (in parentheses). Starting materials in the synthesis examples below are either available from commercial sources such as Aldrich Chemical Co., Milwaukee, Wisconsin, USA, or via literature procedures. Abbreviations used in the examples below have their accepted meanings in the chemical literature. For example, THF (tetrahydrofuran), Et₂O (diethyl ether), MeOH (methanol), CH₂Cl₂ (methylene chloride), LDA (lithium diisopropylamide), MeCN (acetonitrile), and DMAP (4-dimethylaminopyridine).

EXAMPLE 1

This example illustrates preparation of compound 1.



Commercially available 4-(1,1,1,3,3,3-hexafluoro-2-hydroxyisopropyl)aniline (1.93 mmol, 500 mg, Lancaster Synthesis, LTD, Teaneck, New Jersey, USA) was dissolved in THF (8 mL) and triethylamine (3.86 mmol, 391 mg) was added, followed by ethyl isocyanate (1.93 mmol, 137 mg). The mixture was heated at 60°C for 24 hours. The resulting mixture was diluted with CH₂Cl₂ and quenched with water. The organic layer was separated and dried with MgSO₄, the solvent was evaporated and the crude product was recrystallized from MeOH/CH₂Cl₂ to afford 353 mg of 1.


mp 170-172°C.

$^1\text{H-NMR}(\text{CD}_3\text{OD})$: (7.58 (d, $J=8.8$ Hz, 2H), 7.42 (d, $J=8.8$ Hz, 2H), 3.23 (m, 2H), 1.15 (t, $J=7.28$ Hz, 3H). MS (ES+): 331 (M+H, 100). Anal. Calcd. for $\text{C}_{12}\text{H}_{12}\text{N}_2\text{F}_6\text{O}_2$: C, 43.78; H, 3.66; N, 8.48. Found: C, 48.02; H, 3.72; N, 8.36.

5

EXAMPLE 2

This example illustrates preparation of sulfonamide 2.




To a solution of 4-(hexafluoro-2-hydroxyisopropyl)aniline (77.2 mmol, 20 g) in MeOH (200 mL) was added benzenesulfonyl chloride (61.74 mmol, 11.01 g). The mixture was stirred at ambient temperature for 2.5 hours. The solvent was evaporated under reduced pressure and the crude product was dissolved in Et₂O and washed with 3N HCl. The organic layer was separated, the solvent was evaporated and the crude product was purified by silica column chromatography using 25% MeOH/CH₂Cl₂ to afford 14.7g of product.

mp 111-113°C.
 $^1\text{H-NMR}(\text{CDCl}_3)$: δ 7.16 (d, $J=7.16$ Hz, 2H), 7.56 (m, 2H), 7.45 (m, 2H), 7.14 (dd, $J=2.12, 6.84$ Hz, 2H), 6.89 (s, 1H), 3.41 (s, 1H). MS (ES-): 398 (M-H, 100). Anal. Calcd. for $\text{C}_{15}\text{H}_{11}\text{F}_6\text{NO}_3\text{S}$: C, 45.12; H, 2.78; N, 3.51. Found: C, 45.02; H, 2.90; N, 3.54.

The following compounds were prepared in a manner similar to that described above.

25


2.1



mp 205-209°C.

¹H-NMR(d₆-DMSO): δ 11.1 (bs, 1H), 8.61 (s, 1H), 8.08 (d, *J*=8.68 Hz, 1H), 7.89 (d, *J*=8.68 Hz, 1H), 7.55 (d, *J*=8.68 Hz, 2H), 7.20 (dd, *J*=1.96, 6.88 Hz, 2H). MS (ES⁻): 501 (M-H, 100). Anal. Calcd. for C₁₅H₈Cl₃F₆NO₃S: C, 35.84; H, 1.60; N, 2.79. Found: C, 36.42; H, 1.85; N, 2.66.

2.2




2.2

mp 113-115°C.

¹H-NMR(CD₃OD): δ 7.71 (m, 1H), 7.61-7.54 (m, 3H), 7.24 (m, 2H), 7.06 (m, 1H). MS (ES⁺): 404 (M+H⁺, 100). Anal. Calcd. for C₁₃H₉F₆NO₃S: C, 38.52; H, 2.24; N, 3.46. Found: C, 38.79; H, 2.28; N, 3.28.

EXAMPLE 3

This example illustrates the preparation of (3) via N-methylation of (2).



3


To *N*-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]benzenesulfonamide (2) (2.5 g), in DMF (60 mL) was added K₂CO₃ (1.6 g) and methyl iodide (0.64 mL). The reaction was stirred at room temperature for 1.5 hours, then worked up using standard methods, and the crude product was purified by column chromatography to afford the 3.5 g of the title compound.

mp 79-81°C.

¹H-NMR(CDCl₃): δ 7.63-7.45 (m, 7H), 7.26 (m, 2H), 3.50 (s, 3H), 3.22 (s, 3H). MS (ES⁺): 428 (M+H, 100). Anal. Calcd. for C₁₇H₁₅F₆NO₃S: C, 47.78; H, 3.54; N, 3.20; S, 7.50. Found: C, 47.87; H, 3.61; N, 3.27; S, 7.55.

EXAMPLE 4

5 This example illustrates preparation of sulfonamide, 4.




To *N*-methyl-4-(hexafluoro-2-hydroxyisopropyl)aniline (0.915 mmol, 250 mg) prepared as reported by Gilbert *et al.* (*J. Org. Chem.*, **1965**, 30, 1001) in MeOH (2.5 mL) was added *p*-toluenesulfonyl chloride (0.763 mmol, 145 mg). The
10 reaction was stirred overnight at ambient temperature and the solvent was evaporated under reduced pressure. The crude product was dissolved in CH₂Cl₂ and washed with 3N HCl. The organic layer was separated, the solvent evaporated and the crude product was purified by silica column chromatography to afford 130 mg of 4.

mp 98-101°C.

15 ¹H-NMR(CDCl₃): δ 7.64 (d, *J*=8.4 Hz, 2H), 7.39 (d, *J*=8.16 Hz, 2H), 7.21 (m, 4H), 3.16 (s, 3H), 2.40 (s, 3H). MS (ES⁻): 426 (M-H, 100). Anal. Calcd. for C₁₇H₁₅F₆NO₃S: C, 47.78; H, 3.54; N, 3.20; S, 7.50. Found: C, 48.02; H, 3.52; N, 3.20; S, 7.57.

20 The following compounds were prepared using methods similar to those described in this and the preceding examples, with the appropriately substituted arylsulfonyl chlorides or alkylsulfonyl chlorides.

4.1




25

mp 132-134°C.

$^1\text{H-NMR}(\text{CDCl}_3)$: δ 7.64 (d, $J=8.8\text{ Hz}$, 2H), 7.44 (s, 4H), 7.23 (d, $J=8.8\text{ Hz}$, 2H), 3.18 (s, 3H), 1.32 (s, 9H). MS (ES-): 468 (M-H, 100). Anal. Calcd. for $\text{C}_{20}\text{H}_{21}\text{F}_6\text{NO}_3\text{S}$: C, 51.17; H, 4.51; N, 2.98; S, 6.83. Found: C, 51.32; H, 4.47; N, 2.88; S, 6.76.

5

4.2




mp 104-106°C.

$^1\text{H-NMR}(\text{CDCl}_3)$: δ 8.30 (d, $J=8.9\text{ Hz}$, 2H), 7.70 (m, 4H), 7.20 (d, $J=9\text{ Hz}$, 2H), 3.24 (s, 3H). MS (ES-): 457 (M-H, 100). Anal. Calcd. for $\text{C}_{16}\text{H}_{12}\text{F}_6\text{N}_2\text{O}_5\text{S}$: C, 41.93; H, 2.64; N, 6.11; S, 7.00. Found: C, 42.11; H, 2.71; N, 5.95; S, 6.85.

10

4.3




mp 65-69°C.

$^1\text{H-NMR}(\text{CDCl}_3)$: δ 7.63 (d, $J=8.7\text{ Hz}$, 2H), 7.34 (m, 3H), 7.03 (dd, 8.9, 3.0 Hz, 1H), 6.85 (d, $J=9.04\text{ Hz}$, 1H), 3.74 (s, 3H), 3.54 (s, 3H), 3.32 (s, 3H). MS (ES+): 474 (M+H, 100). Anal. Calcd. for $\text{C}_{18}\text{H}_{17}\text{F}_6\text{NO}_5\text{S}$: C, 45.67; H, 3.62; N, 2.96; S, 6.77. Found: C, 45.21; H, 3.71; N, 2.93; S, 6.58.

15

20


4.4



mp 109-111°C.

¹H-NMR(CDCl₃): δ 8.4 (d, *J*=8.4 Hz, 2H), 7.44 (d, *J*=8.8 Hz, 2H), 7.21 (d, *J*=8.9 Hz, 2H), 6.90 (d, *J*=8.9 Hz, 2H), 3.84 (s, 3H), 3.16 (s, 3H). MS (ES⁺): 444 (M+H, 100). Anal. Calcd. for C₁₇H₁₅F₆NO₄S: C, 46.05; H, 3.41; N, 3.16; S, 7.23. Found: C, 45.84; H, 3.41; N, 3.12; S, 7.14.


4.5



mp 175-180°C.

¹H-NMR(CDCl₃): δ 8.21 (d, *J*=8.8 Hz, 1H), 8.12 (d, *J*=7.6 Hz, 1H), 7.64 (dd, *J*=7.08, 8.8 Hz, 1H), 7.56 (d, *J*=8.76 Hz, 2H), 7.21 (d, *J*=8.5 Hz, 2H), 3.53 (s, 3H). MS (ES⁻): 470 (M-H, 100). Anal. Calcd. for C₁₆H₁₁F₆N₃O₃S₂: C, 40.77; H, 2.35; N, 8.91; S, 13.60. Found: C, 40.64; H, 2.22; N, 8.84; S, 13.78.


4.6



mp 113-115°C.

¹H-NMR(CDCl₃): δ 8.06 (dd, *J*=9.12, 0.7 Hz, 1H), 7.85 (dd, *J*=6.88, 0.8 Hz, 1H), 7.60 (d, *J*=9.0 Hz, 2H), 7.46 (dd, *J*=9.08, 6.84 Hz, 1H), 7.25 (d, *J*=9.0 Hz, 2H), 3.55 (s, 3H).


4.7



mp 195-197°C.

- 5 $^1\text{H-NMR}(\text{d}_6\text{-DMSO})$: δ 8.76 (s, 1H), 7.77 (d, $J=2.5$ Hz, 1H), 7.64 (d, $J=8.44$ Hz, 2H), 7.56 (dd, $J=8.8, 2.5$ Hz, 1H), 7.30 (d, $J=8.9$ Hz, 2H), 7.07 (d, $J=9.0$ Hz, 1H), 3.13 (s, 3H).

4.8




10

mp 90-91°C.

- $^1\text{H-NMR}(\text{CDCl}_3)$: δ 7.66 (d, $J=8.5$ Hz, 2H), 7.58 (d, $J=8.6$ Hz, 2H), 7.37 (d, $J=8.64$ Hz, 2H), 7.23 (d, $J=8.8$ Hz, 2H), 3.19 (s, 3H). MS (ES⁻): 491 (M-H, 100), 489 (M-H, 100). Anal. Calcd. for $\text{C}_{16}\text{H}_{12}\text{BrF}_6\text{NO}_3\text{S}$: C, 39.04; H, 2.46; N, 2.85; S, 6.51; Br, 16.23. Found: C, 39.24; H, 2.44; N, 2.71; S, 6.52; Br, 16.38.
- 15


4.9



mp 98-99°C.

$^1\text{H-NMR}(\text{CDCl}_3)$: δ 7.67 (d, $J=9$ Hz, 2H), 7.54 (m, 2H), 7.21 (d, $J=9.0$ Hz, 2H), 7.13 (m, 2H), 3.20 (s, 3H). MS (ES-): 430 (M-H, 100). Anal. Calcd. for $\text{C}_{16}\text{H}_{12}\text{F}_7\text{NO}_3\text{S}$: C, 44.55; H, 2.80; N, 3.25; S, 7.43. Found: C, 44.54; H, 2.83; N, 3.18; S, 7.39.


5

4.10

mp 98-100°C.

$^1\text{H-NMR}(\text{CDCl}_3)$: δ 8.8 (d, $J=8.8$ Hz, 2H), 7.55 (m, 1H), 7.49 (s, 1H), 7.38 (m, 2H), 7.21 (d, $J=9$ Hz, 2H), 3.21 (s, 3H). MS (ES-): 446 (M-H, 100). Anal. Calcd. for $\text{C}_{16}\text{H}_{12}\text{ClF}_7\text{NO}_3\text{S}$: C, 42.92; H, 2.70; N, 3.13; S, 7.16; Cl, 7.92. Found: C, 42.90; H, 2.62; N, 3.02; S, 7.15; Cl, 8.04.

10

4.11


15

mp 114-116°C.

$^1\text{H-NMR}(\text{CDCl}_3)$: δ 7.94 (d, $J=6.5$ Hz, 1H), 7.62 (d, $J=8.8$ Hz, 2H), 7.48 (m, 2H), 7.33 (m, 3H), 3.42 (s, 3H). MS (ES-): 446 (M-H, 100). Anal. Calcd. for $\text{C}_{16}\text{H}_{12}\text{ClF}_7\text{NO}_3\text{S}$: C, 42.92; H, 2.70; N, 3.13; S, 7.16; Cl, 7.92. Found: C, 42.84; H, 2.62; N, 3.02; S, 7.24; Cl, 8.03.

20

4.12




4.12

mp 124-126°C.

¹H-NMR(CDCl₃): δ 7.68 (m, 4H), 7.42 (m, 1H), 7.33 (m, 1H), 7.21 (d, *J*=8.08 Hz, 2H), 3.21 (s, 3H). Anal. Calcd. for C₁₆H₁₂BrF₆NO₃S: C, 39.04; H, 2.46; N, 2.85; S, 6.51; Br, 16.23. Found: C, 39.04; H, 2.38; N, 2.75; S, 6.45; Br, 16.36.

4.13




4.13

mp 79-81°C.

¹H-NMR(CDCl₃): δ 7.83 (d, *J*=7.0 Hz, 1H), 7.74 (d, *J*=7.0 Hz, 1H), 7.66 (m, 3H), 7.63 (m, 1H), 7.18 (d, *J*=7.0 Hz, 2H), 3.20 (s, 3H). MS (ES⁻): 480 (M-H, 100). Anal. Calcd. for C₁₇H₁₂F₉NO₃S: C, 42.42; H, 2.51; N, 3.91; S, 6.66. Found: C, 42.66; H, 2.48; N, 2.96; S, 6.49.

4.14




4.14

mp 135-137°C.

MS (ES⁻): 430 (M-H, 100). Anal. Calcd. for C₁₆H₁₂F₇NO₃S: C, 44.55; H, 2.80; N, 3.25; S, 7.43. Found: C, 44.45; H, 2.76; N, 3.17; S, 7.48.

4.15




4.15

mp 135-137°C.

MS (ES⁻): 430 (M-H, 100). Anal. Calcd. for C₁₆H₁₂F₇NO₃S: C, 44.55;
 5 H, 2.80; N, 3.25; S, 7.43. Found: C, 44.65; H, 2.84; N, 3.17; S, 7.39.

4.16




4.16

mp 115-118°C.

10 ¹H-NMR(CDCl₃): δ 7.74-7.65 (m, 6H), 7.22 (d, J=6.72 Hz, 2H), 3.22 (s, 3H). MS (ES⁻): 430 (M-H, 100). Anal. Calcd. for C₁₇H₁₂F₉NO₃S: C, 42.42; H, 2.51; N, 3.91; S, 6.66. Found: C, 42.24; H, 2.42; N, 2.87; S, 6.65.

4.17




4.17

mp 88-89°C.


15 ¹H-NMR(d₆-DMSO): δ 8.80 (s, 1H), 7.75-7.50 (m, 7H), 7.25 (m, 2H), 3.60 (m, 2H), 3.31 (s, 3H), 0.977 (m, 3H). MS (ES⁺): 428 (M+H, 100).

4.18



¹H-NMR(CD₃OD): δ 8.57 (dd, *J*=1.12, 8.56 Hz, 1H), 8.13 (m, 1H), 8.36 (m, 1H), 7.61-7.52 (m, 3H), 7.23-7.15 (m, 4H), 3.22 (s, 3H), 2.84 (s, 6H). MS (ES⁺): 507 (M+H, 100).


4.19



mp 180-183°C.

¹H-NMR(CD₃OD): δ 8.92 (dd, *J*=4.2, 1.76 Hz, 1H), 8.39 (dd, *J*=8.36, 1.72 Hz, 1H), 8.35 (dd, *J*=1.36, 7.36 Hz, 1H), 8.17 (dd, *J*=1.32, 8.2 Hz, 1H), 7.63 (m, 2H), 7.52 (m, 2H), 7.25 (dd, *J*=6.88, 2.0 Hz, 2H), 3.61 (s, 3H). MS (ES⁺): 465 (M+H, 100).


4.20



mp 208-210°C.

¹H-NMR(CD₃OD): δ 8.12 (d, *J*=8.56 Hz, 2H), 7.70 (d, *J*=8.72 Hz, 2H), 7.64 (d, *J*=8.6 Hz, 2H), 7.26 (d, *J*=8.96 Hz, 2H), 3.24 (s, 3H). MS (ES⁻): 456 (M-H, 100).


4.21



mp 102-106°C.

¹H-NMR(CD₃OD): δ 7.72 (d, *J*=8.4 Hz, 2H), 7.36 (dd, *J*=2.12, 6.88 Hz, 2H), 3.31 (s, 3H), 2.22 (s, 3H), 1.86 (s, 3H). MS (ES⁺): 492 (M+H, 100).


4.22



mp 204-206°C.

¹H-NMR(CD₃OD): δ 9.30 (s, 1H), 8.41 (m, 2H), 8.13 (d, *J*=1.5 Hz, 1H), 7.83 (t, *J*=8.03 Hz, 1H), 7.74 (d, *J*=6.32 Hz, 1H), 7.60 (d, *J*=8.8 Hz, 2H), 7.20 (dd, *J*=2.08, 6.84 Hz, 2H), 3.24 (s, 3H). MS (ES⁺): 465 (M+H, 50).


4.23



mp 199-202°C.

¹H-NMR(CD₃OD): δ 8.26 (m, 1H), 7.71 (m, 3H), 7.63 (t, *J*=8.12 Hz, 1H), 7.25 (dd, *J*=2.12, 6.8 Hz, 2H), 3.23 (s, 3H). MS (ES⁺): 458 (M+H, 100).

4.24




4.24

mp 133-136°C.

¹H-NMR(CD₃OD): δ 7.82 (dd, *J*=5.08, 1.28 Hz, 1H), 7.70 (d, *J*=8.52 Hz, 2H), 7.38 (dd, *J*=1.2, 3.74 Hz, 1H), 7.28 (m, 2H), 7.16 (m, 1H), 3.26 (s, 3H). MS (ES⁺): 420 (M+H, 100).

4.25




4.25

10

mp 120-122°C.

¹H-NMR(CD₃OD): δ 7.68 (d, *J*=8.72 Hz, 2H), 7.61 (s, 1H), 7.43 (s, 1H), 7.38 (m, 2H), 3.40 (s, 3H). MS (ES⁻): 487 (M-H, 100).

4.26




4.26

15

Oil.

¹H-NMR(CD₃OD): δ 7.90 (m, 1H), 7.70-7.63 (m, 3H), 7.40 (m, 2H), 3.42 (s, 3H). MS (ES⁻): 515 (M-H, 100).


4.27



mp 148-149°C.

¹H-NMR(CD₃OD): δ 8.80 (d, *J*=4.76 Hz, 1H), 8.64 (s, 1H), 7.96 (m, 1H), 7.72 (d, *J*=8.72 Hz, 2H), 7.59 (m, 1H), 7.28 (d, *J*=9 Hz, 2H), 3.26 (s, 3H). MS (ES⁻): 413 (M-H, 100).


4.28



mp 130-134°C.

¹H-NMR(CD₃OD): δ 7.64 (m, 1H), 7.54 (m, 6H), 7.12 (dd, *J*=6.64, 2.12 Hz, 2H), 3.18 (s, 3H). MS (ES⁺): 459 (M+H⁺, 100). Anal. Calcd. for C₁₆H₁₂F₆N₂O₅S: C, 41.93; H, 2.64; N, 6.11. Found: C, 41.64; H, 2.69; N, 6.10.


4.29



mp 83-86°C.

¹H-NMR(CD₃OD): δ 7.82 (m, 1H), 7.73 (m, 2H), 7.40 (m, 1H), 7.24 (m, 2H), 7.15 (m, 1H), 3.71 (q, *J*=7.16, 14.24 Hz, 2H), 1.09 (t, *J*= 7.24 Hz, 3H). MS (ES⁺): 434 (M+H⁺, 100).

4.30




4.30

mp 87-89°C.

- 5 ¹H-NMR(CD₃OD): δ 8.86 (s, 1H), 8.07 (dd, *J*=1.28, 4.92 Hz, 1H), 7.71 (m, 2H), 7.36 (m, 2H), 7.28 (m, *J*=3.84, 4.98 Hz, 1H), 4.61 (q, *J*=8.8, 17.6 Hz, 2H).

4.31




4.31

- 10 mp 142-145°C.

¹H-NMR(CD₃OD): δ 7.68 (m, 5H), 7.44 (s, 1H), 7.28 (m, 2H), 3.24 (s, 3H), 3.06 (s, 3H), 2.84 (s, 3H). Anal. Calcd. for C₁₉H₁₈F₆N₂O₄S: C, 47.11; H, 3.75; N, 5.78. Found: C, 47.19; H, 3.75; N, 5.77.

15


4.32



4.32

- ¹H NMR (CDCl₃): δ 7.70 (d, *J* = 8.4 Hz, 2H), 7.50 (m, 4H), 7.25 (d, *J* = 8.4 Hz, 2H), 6.78 (dd, *J* = 16.0, 11.0 Hz, 1H), 5.90 (d, *J* = 16.0 Hz, 1H), 5.48 (d, *J* = 11.0 Hz, 1H), 3.55 (s, 1H), 3.28 (s, 3H). MS (ES⁻): 438 (M-H, 100).
- 20


4.33



4.33

¹H NMR (CDCl₃): δ 7.70 (d, *J* = 8.5 Hz, 2H), 7.42 (d, *J* = 5.0 Hz, 1H),
 5 7.35 (d, *J* = 8.5 Hz, 2H), 7.25 (d, *J* = 5 Hz, 1H), 3.75 (s, 3H), 3.65 (s, 1H), 3.35 (s, 3H). MS (ES⁺): 478 (M+H, 100).

4.34




4.34

10

¹H NMR (CDCl₃): δ 7.91 (d, *J* = 7.5 Hz, 1H), 7.68 (d, *J* = 8.4 Hz, 2H),
 7.63 (t, *J* = 8.4 Hz, 1H), 7.36 (t, *J* = 7.5 Hz, 2H), 7.32 (d, *J* = 8.4 Hz, 2H), 3.60 (s, 1H), 3.30 (s, 3H). MS (ES⁺): 498 (M+H, 100).

4.35




4.35

15

¹H NMR (CDCl₃): δ 7.65 (d, *J* = 8.2 Hz, 2H), 7.28 (d, *J* = 8.2 Hz, 2H),
 6.95 (d, *J* = 5.0 Hz, 1H), 6.70 (d, *J* = 5.0 Hz, 1H), 3.82 (s, 1H), 3.35 (s, 3H). MS (ES-
 20): 492 (M-H, 100).

4.36




4.36

^1H NMR (CDCl_3): δ 8.05 (d, $J = 7.5$ Hz, 2H), 7.76 (d, $J = 7.5$ Hz, 2H), 7.70 (d, $J = 8.2$ Hz, 2H), 7.21 (d, $J = 8.2$ Hz, 2H), 3.57 (s, 1H), 3.21 (s, 3H), 3.06 (s, 3H). MS (ES $^-$): 490 (M-H, 100).

5

4.37




4.37

^1H NMR (CDCl_3): δ 7.68 (d, $J = 8.5$ Hz, 2H), 7.50 (s, 1H), 7.40 (d, $J = 8.5$ Hz, 2H), 7.20 (s, 1H), 3.70 (s, 3H), 3.30 (s, 3H). MS (ES $^+$): 418 (M+H, 100).

10

4.38




4.38

^1H NMR (CD_3OD): δ 7.70 (d, $J = 8.5$ Hz, 2H), 7.43 (d, $J = 8.5$ Hz, 2H), 7.33 (d, $J = 5.0$ Hz, 1H), 6.92 (d, $J = 5.0$ Hz, 1H), 3.46 (s, 2H). MS (ES $^-$): 925 (2M-H, 100). Anal. Calcd. for $\text{C}_{15}\text{H}_{11}\text{F}_6\text{NO}_5\text{S}_2$: C, 38.88; H, 2.39; N, 3.02; S, 13.84. Found: C, 39.65; H, 2.97; N, 2.70; S, 12.09.

15

4.39




4.39

^1H NMR (CDCl_3): δ 7.90 (d, $J = 7.5$ Hz, 1H), 7.78 (d, $J = 7.5$ Hz, 1H), 7.70 (t, $J = 7.5$ Hz, 1H), 7.68 (d, $J = 8.4$ Hz, 2H), 7.57 (t, $J = 7.5$ Hz, 1H), 7.27 (d, $J = 8.4$ Hz, 2H), 3.83 (s, 1H), 3.36 (s, 3H). MS (ES $^-$): 480 (M-H, 100). Anal. Calcd. for $\text{C}_{17}\text{H}_{12}\text{F}_9\text{NO}_3\text{S}$: C, 42.42; H, 2.51; N, 2.91; S, 6.66. Found: C, 43.71; H, 2.75; N, 2.67; S, 6.92.

20

25


4.40



4.40

¹H NMR (CDCl₃): δ 7.72 (d, *J* = 7.2 Hz, 2H), 7.69 (d, *J* = 8.2 Hz, 2H),
 5 7.66 (d, *J* = 7.2 Hz, 2H), 7.21 (d, *J* = 8.2 Hz, 2H), 3.52 (s, 1H), 3.20 (s, 3H). MS (ES-): 480 (M-H, 100).


4.41



4.41

¹H NMR (CDCl₃): δ 7.72 (d, *J* = 7.2 Hz, 2H), 7.35 (d, *J* = 8.4 Hz, 2H),
 10 3.58 (s, 1H), 3.25 (s, 3H), 2.25 (s, 3H), 1.96 (s, 3H). MS (ES-): 431 (M-H, 100).


4.42



4.42

¹H NMR (CDCl₃): δ 8.56 (s, 1H), 7.78 (t, *J* = 7.2 Hz, 1H), 7.65 (d, *J* =
 15 8.4 Hz, 2H), 7.50 (d, *J* = 4.5 Hz, 1H), 7.30 (d, *J* = 8.4 Hz, 2H), 7.28 (m, 2H), 7.21 (d,
 20 *J* = 4.5 Hz, 1H), 3.30 (s, 3H). MS (ES-): 495 (M-H, 100).

4.43




4.43

^1H NMR (CDCl_3): δ 7.72 (d, $J = 7.5$ Hz, 1H), 7.65 (d, $J = 8.4$ Hz, 2H), 7.50 (s, 1H), 7.48 (d, $J = 7.5$ Hz, 1H), 7.29 (d, $J = 8.4$ Hz, 2H), 3.85 (s, 1H), 3.35 (s, 3H). MS (ES $^-$): 576 (M-H, 100).

5

4.44




4.44

^1H NMR (CDCl_3): δ 7.86 (d, $J = 7.8$ Hz, 1H), 7.85 (s, 1H), 7.70 (d, $J = 8.5$ Hz, 2H), 7.60 (d, $J = 7.8$ Hz, 2H), 7.20 (d, $J = 8.5$ Hz, 2H), 3.48 (s, 1H), 3.25 (s, 3H). MS (ES $^-$): 437 (M-H, 100). Anal. Calcd. for $\text{C}_{17}\text{H}_{12}\text{F}_6\text{N}_2\text{O}_3\text{S}$: C, 46.58; H, 2.76; N, 6.39; S, 7.32. Found: C, 46.97; H, 2.92; N, 6.18; S, 7.17.

10

4.45




4.45

15

^1H NMR (CDCl_3): δ 7.88 (d, $J = 8.0$ Hz, 1H), 7.65 (d, $J = 8.0$ Hz, 1H), 7.64 (d, $J = 8.4$ Hz, 2H), 7.35 (d, $J = 8.4$ Hz, 2H), 7.26 (t, $J = 8.0$ Hz, 1H), 3.45 (s, 3H), 3.44 (s, 1H). MS (ES $^-$): 481 (M-H, 100). Anal. Calcd. for $\text{C}_{16}\text{H}_{11}\text{Cl}_2\text{F}_6\text{NO}_3\text{S}$: C, 39.85; H, 2.30; Cl, 14.70; N, 2.90; S, 6.65. Found: C, 40.49; H, 2.37; N, 2.89; S, 6.60; Cl, 14.95.

20

4.46




4.46

25

^1H NMR (CDCl_3): δ 7.71 (d, $J = 8.4$ Hz, 2H), 7.35 (d, $J = 8.5$ Hz, 2H), 6.86 (s, 1H), 3.41 (s, 1H), 3.35 (s, 3H). MS (ES $^-$): 486 (M-H, 100). Anal. Calcd. for

$C_{14}H_9C_{12}F_6NO_3S_2$: C, 34.44; H, 1.86; Cl, 14.52; N, 2.87; S, 13.13. Found: C, 35.20; H, 1.87; N, 2.95; S, 13.31; Cl, 15.04.


4.47



4.47

1H NMR ($CDCl_3$): δ 7.75 (d, J = 8.4 Hz, 2H), 7.48 (d, J = 8.4 Hz, 2H), 3.49 (s, 1H), 3.37 (s, 3H), 2.88 (s, 3H). MS (ES⁻): 350 (M-H, 100). Anal. Calcd. for $C_{11}H_{11}F_6NO_3S$: C, 37.61; H, 3.16; N, 3.99; S, 9.13. Found: C, 37.83; H, 3.27; N, 4.03; S, 9.28.


4.48



4.48

1H NMR ($CDCl_3$): δ 7.70 (d, J = 8.0 Hz, 2H), 7.49 (d, J = 8.0 Hz, 2H), 3.45 (s, 1H), 3.30 (s, 3H), 2.80 (s, 6H). MS (ES⁻): 379 (M-H, 100). Anal. Calcd. for $C_{12}H_{14}F_6N_2O_3S$: C, 37.90; H, 3.71; N, 7.37. Found: C, 38.05; H, 3.77; N, 7.45.


4.49



4.49

1H NMR ($CDCl_3$): δ 7.72 (d, J = 8.2 Hz, 2H), 7.48 (d, J = 8.2 Hz, 2H), 3.65 (s, 1H), 3.30 (s, 3H), 3.01 (t, J = 6.3 Hz, 2H), 1.80 (m, 2H), 1.31 (m, 2H), 0.92 (t, J = 6.0 Hz, 3H). MS (ES⁻): 392 (M-H, 100).


4.50



4.50

¹H NMR (CDCl₃): δ 7.71 (d, *J* = 8.2 Hz, 2H), 7.50 (d, *J* = 8.2 Hz, 2H),
 5 3.90 (s, 1H), 3.41 (s, 3H), 3.32 (m, 1H), 1.38 (d, *J* = 6.0 Hz, 6H). MS (ES⁻): 378 (M-H, 100).

4.51




4.51

10

¹H NMR (CDCl₃): δ 7.60 (d, *J* = 8.0 Hz, 2H), 7.28 (d, *J* = 8.0 Hz, 2H),
 7.25 (d, *J* = 2.5 Hz, 1H), 7.15 (dd, *J* = 7., 2.5 Hz, 1H), 7.06 (d, *J* = 7.2 Hz, 1H), 4.30
 (q, *J* = 7.0 Hz, 2H), 4.20 (q, *J* = 7.0 Hz, 2H), 3.50 (s, 1H), 3.38 (s, 3H). MS (ES⁻): 610
 (M+H, 100).

15


4.52



4.52

¹H NMR (CDCl₃): δ 7.78 (d, *J* = 7.0 Hz, 2H), 7.70 (d, *J* = 8.2 Hz, 2H),
 20 7.62 (d, *J* = 7.0 Hz, 2H), 7.20 (d, *J* = 8.2 Hz, 2H), 3.51 (s, 1H), 3.21 (s, 3H). MS (ES⁻): 437 (M-H, 100).

4.53




4.53

^1H NMR (CDCl_3): δ 7.82 (m, 2H), 7.68 (d, $J = 8.4$ Hz, 2H), 7.65 (m, 2H), 7.28 (d, $J = 8.4$ Hz, 2H), 3.72 (s, 1H), 3.40 (s, 3H). MS (ES $^-$): 437 (M-H, 100).

5

4.54




4.54

^1H NMR (CDCl_3): δ 7.90 (s, 1H), 7.66 (d, $J = 8.4$ Hz, 2H), 7.42 (m, 2H), 7.33 (d, $J = 8.4$ Hz, 2H), 3.47 (s, 1H), 3.40 (s, 3H). MS (ES $^-$): 481 (M-H, 100).

10 Anal. Calcd. for $\text{C}_{16}\text{H}_{11}\text{Cl}_2\text{F}_6\text{NO}_3\text{S}$: C, 39.85; H, 2.30; N, 2.90; S, 6.65. Found: C, 40.01; H, 2.19; N, 2.99; S, 6.82.

4.55




15

4.55

^1H NMR (CDCl_3): δ 7.80 (s, 1H), 7.77 (d, $J = 8.4$ Hz, 2H), 7.31 (d, $J = 8.4$ Hz, 2H), 3.50 (s, 1H), 3.31 (s, 3H). MS (ES $^-$): 496 (M-H, 100).

4.56




20

4.56

^1H NMR (CDCl_3): δ 7.72 (d, $J = 8.4$ Hz, 2H), 7.30 (d, $J = 8.4$ Hz, 2H), 7.12 (s, 1H), 3.48 (s, 1H), 3.28 (s, 3H). MS (ES $^-$): 531 (M+H, 100). Anal. Calcd. for $\text{C}_{14}\text{H}_9\text{BrClF}_6\text{NO}_3\text{S}_2$: C, 31.57; H, 1.70; N, 2.63; S, 12.21. Found: C, 40.01; H, 2.19; N, 2.99; S, 6.82.

25


4.57



4.57

¹H NMR (CDCl₃): δ 7.66 (d, *J* = 8.4 Hz, 2H), 7.35 (, 3H), 7.20 (m, 3H),
 5 3.50 (s, 1H), 3.19 (s, 3H), 2.30 (s, 3H). MS (ES⁻): 426 (M-H, 100).

4.58




4.58

¹H NMR (CD₃OD): δ 8.59 (s, 1H), 7.97 (2H, *J* = 7.8 Hz, 2H), 7.70 (t, *J* = 7.8 Hz, 1H), 7.68 (d, *J* = 8.2 Hz, 2H), 7.62 (t, *J* = 7.8 Hz, 2H), 7.50 (s, 1H), 7.20 (d, *J* = 8.2 Hz, 2H), 3.30 (s, 1H), 3.21 (s, 3H). MS (ES⁻): 558 (M-H, 100). Anal. Calcd. for C₂₀H₁₅F₆NO₅S₃: C, 42.93; H, 2.73; N, 2.50; S, 17.19. Found: C, 43.07, H, 2.70; N, 2.50; S, 17.30.

15

4.59




4.59

This compound was prepared from 4.46 using conditions similar to the
 20 palladium-catalyzed reduction of Example 5, below (monitoring the reaction to obtain an incomplete reduction of the chloro substituents).

¹H NMR (CDCl₃): δ 7.70 (d, *J* = 8.8 Hz, 2H), 7.50 (s, 1H), 7.28 (d, *J* = 8.8 Hz, 2H), 6.83 (s, 1H), 3.37 (s, 1H), 3.25 (s, 1H). MS (ES⁻): 907 (2M-H, 100).
 Anal. Calcd. for C₁₄H₁₀ClF₆NO₃S₂: C, 37.05; H, 2.22; N, 3.09. Found: C, 37.92; H,
 25 2.35; N, 3.11.

4.60




4.60

This compound was prepared from 4.46 using conditions similar to the
 5 palladium-catalyzed reduction of Example 5, below.

¹H NMR (CDCl₃): δ 7.76 (d, *J* = 3.0 Hz, 1H), 7.67 (d, *J* = 8.8 Hz, 2H),
 7.37 (dd, *J* = 4.0, 1.5 Hz, 1H), 7.27 (d, *J* = 8.8 Hz, 2H), 6.99 (d, *J* = 4.0 Hz, 1H), 3.49
 (s, 1H), 3.22 (s, 3H). MS (ES⁻): 461 (M-H, 100). MS (ES⁻): 837 (2M-H, 100). Anal.
 Calcd. for C₁₄H₁₁F₆NO₃S₂: C, 40.10; H, 2.64; N, 3.34. Found: C, 40.22; H, 2.67; N,
 10 3.36.

4.61



4.61


The starting aniline was prepared via treatment of 4-
 15 (trifluoroacetyl)aniline with methyl magnesium bromide. Subsequent alkylation and
 sulfonylation was carried out as described in the examples above.

¹H-NMR(CD₃OD): δ 7.64 (m, 1H), 7.54 (m, 6H), 7.12 (dd, *J*=6.64,
 2.12, 2H), 3.18 (s, 3H). MS (ES⁺): 382 (M+Na⁺, 50). Oil

20

EXAMPLE 5

This example illustrates preparation of compound 5 by reduction of
 4.28.



5


25

To a mixture of *N*-Methyl-*N*-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]-3-nitrobenzenesulfonamide (**4.28**) (70 mg), dissolved in 10 mL of MeOH was added 10% Pd/carbon (10mg) and the reaction was stirred under a H₂ atmosphere for 2 days. The resulting mixture was filtered through Celite and the solvent was evaporated under reduced pressure. The crude product was

purified by preparative-TLC to obtain 30 mg of the title compound as an oil.
¹H-NMR(CD₃OD): δ 7.66 (d, *J*=8.76 Hz, 2H), 7.23 (m, 3H), 6.91 (d, *J*=7.9 Hz, 1H), 6.86 (d, *J*=1.62, 7.96 Hz, 1H), 6.74 (s, 1H), 3.19 (s, 3H). Anal. Calcd. For C₁₉H₁₈F₆N₂O₄S: C, 47.11; H, 3.75; N, 5.78. Found: C, 47.19; H, 3.75; N, 5.77.

EXAMPLE 6

This example illustrates preparation of compound **6** via esterification of **4.23**.




To *N*-Methyl-*N*-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]-3-carboxybenzenesulfonamide (**4.23**, 0.536 mmol, 245 mg) under N₂ at 0°C, in 10 mL of THF was added triethylamine (2.15 mmol, 300 μL), MeOH (0.864 mmol, 35 μL), and bis(2-oxo-3-oxazolidinyl)phosphinic chloride (0.833 mmol, 212 mg). The mixture was stirred for 20 h at 0°C. The reaction was quenched with water, the THF was removed under vacuum, and the reaction product was extracted with EtOAc. The organic extract was dried with MgSO₄, the solvent was evaporated and the crude product was purified by preparative TLC to afford 106 mg of the title compound as a solid.

mp 88-90°C.

¹H-NMR(CD₃OD): δ 8.26 (d, *J*=7.68 Hz, 1H), 8.03 (d, *J*=1.7 Hz, 1), 7.73 (m, 3H), 7.68 (m, 3H), 7.25 (m, 2H) 3.89 (s, 3H), 3.22 (s, 3H). MS (ES⁺): 472 (M+H⁺, 100). Anal. Calcd. for C₁₈H₁₅F₆NO₅S: C, 45.86; H, 3.21; N, 2.97. Found: C, 46.02; H, 3.32; N, 2.91.

EXAMPLE 7

This example provides the synthesis of compound 7.




7

To a stirred solution of N-methyl-4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]aniline (80 mg, 0.293 mmol) and diisopropylethylamine (112 mL, 0.630 mmol) in dichloromethane (2 mL) was added isobutyl chloroformate (42 mL, 0.323 mmol) at 0°C. The resulting mixture was stirred for 6 h as the cooling bath temperature increased to room temperature. The reaction mixture was diluted with ethyl acetate, and the organic phase was washed sequentially with 1N HCl and brine. The separated organic layer was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography on silica gel (hexanes and ethyl acetate (4:1) as eluant) to give the title compound (80 mg, 73.2%) as a clear oil.

¹H NMR (CDCl₃): δ 7.70 (d, *J*=8.5 Hz, 2H), 7.39 (d, *J*=8.5 Hz, 2H), 3.93 (d, *J*=7.2 Hz, 2H), 3.69 (s, 1H), 3.37 (s, 3H), 1.92 (m, 1H), 0.90 (d, *J*= 6.3 Hz, 6H). Anal. Calcd. for C₁₅H₁₇F₆NO₃: C, 48.26; H, 4.59; N, 3.75. Found: C, 48.34; H, 4.61; N, 3.72.


The following carbamates were prepared according to the procedure described.

7.1




¹H NMR (CDCl₃): δ 7.71 (d, *J*=8.6 Hz, 2H), 7.43 (d, *J*=8.6 Hz, 2H), 7.05 (m, 4H), 4.46 (s, 1H), 3.42 (s, 3H). MS (ES⁻): 412 (M+H, 100).

7.2




¹H NMR (CDCl₃): δ 7.69 (d, *J*=8.4 Hz, 2H), 7.37 (d, *J*=8.4 Hz, 2H), 5.75 (m, 1H), 5.05 (d, *J*=16.5 Hz, 1H), 5.04 (d, *J*=10.0 Hz, 1H), 4.20 (t, *J*=6.5 Hz, 2H), 3.40 (s, 1H), 3.35 (s, 3H), 2.40 (m, 2H). MS (ES⁻): 370 (M-H, 100).

7.3




¹H NMR (CDCl₃): δ 7.69 (d, *J*=8.4 Hz, 2H), 7.35 (d, *J*=8.4 Hz, 2H), 5.15 (s, 1H), 4.30 (m, 2H), 3.60 (m, 2H). MS (ES⁻): 374 (M-H, 100).

7.4




¹H NMR (CDCl₃): δ 8.05 (d, *J*=7.8 Hz, 2H), 7.75 (d, *J*=8.5 Hz, 2H), 7.43 (d, *J*=8.5 Hz, 2H), 7.20 (d, *J*=7.8 Hz, 2H), 4.81 (s, 1H), 3.90 (s, 3H), 3.45 (s, 3H). MS (ES⁻): 452 (M+H, 100).

7.5



¹H NMR (CDCl₃): δ 7.70 (d, *J*=8.6 Hz, 2H), 7.37 (d, *J*=8.6 Hz, 2H), 3.81 (s, 2H), 1.10 (s, 9H). MS (ES⁻): 386 (M-H, 100). Anal. Calcd. for C₁₇H₁₃F₆NO₃: C, 49.62; H, 4.94; N, 3.62. Found: C, 49.78; H, 5.02; N, 3.64.

7.6




¹H NMR (CDCl₃): δ 7.70 (d, *J*=8.6 Hz, 2H), 7.35 (d, *J*=8.6 Hz, 2H), 7.30 (m, 5H), 5.18 (s, 2H), 5.05 (s, 1H), 3.32 (s, 3H). MS (ES⁻): 408 (M-H, 100).

5

EXAMPLE 8

This example provides the synthesis of urea (8).



10

8

A mixture of N-methyl-4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]aniline (50 mg, 0.83 mmol), N,N-dimethylaminopyridine (DMAP, 22 mg, 0.183 mmol) and 4-trifluoromethylphenylisocyanate (26 mL, 0.183 mmol) in toluene (1 mL) was stirred at room temperature until reaction was completed by tlc. The resulting mixture was diluted with ethyl acetate, washed with 1N HCl and brine. The organic layer was dried over MgSO₄, and filtered. Solvents were evaporated and the residue was purified by flash chromatography on silica gel (hexanes:ethyl acetate (4:1) as eluant) to provide the title compound (8, 70.0 mg, 85%).

15


¹H NMR (DMSO): δ 9.00 (s, 1H), 7.69 (d, *J*=8.2 Hz, 4H), 7.48 (d, *J*=8.2 Hz, 2H), 7.48 (d, *J*=8.2 Hz, 2H), 3.35 (s, 3H). MS (ES⁺): 459 (M-H, 100).

20

The following ureas were prepared as described above, using the appropriate isocyanates:

25


8.1



$^1\text{H NMR}$ (CD_3OD): δ 7.80 (d, $J=8.2$ Hz, 2H), 7.45 (d, $J=8.2$ Hz, 2H), 7.30 (d, $J=8.0$ Hz, 2H), 6.98 (t, 8.0 Hz, 2H), 3.35 (s, 3H).

5


8.2



$^1\text{H NMR}$ (CD_3OD): δ 7.80 (d, $J=8.2$ Hz, 2H), 7.45 (d, $J=8.5$ Hz, 2H), 7.20 (d, $J=8.0$ Hz, 2H), 6.82 (d, $J=8.0$ Hz, 2H), 4.85 (s, 1H), 3.72 (s, 3H), 3.35 (s, 3H), 3.38 (s, 3H). MS (ES $^-$): 421 (M+H, 100).

10

8.3




$^1\text{H NMR}$ (CDCl_3): δ 7.81 (d, $J=8.4$ Hz, 2H), 7.42 (d, $J=8.4$ Hz, 2H), 7.20-7.32 (m, 4H), 7.00 (t, $J=7.0$ Hz, 1H), 6.25 (bs, 1H), 3.91 (bs, 1H), 3.38 (s, 3H). MS (ES $^-$): 193 (M-H, 100). Anal. Calcd. for $\text{C}_{17}\text{H}_{14}\text{F}_6\text{N}_2\text{O}_2$: C, 52.05; H, 3.60; N, 7.14. Found: C, 52.09; H, 3.70; N, 7.29.

15

20

EXAMPLE 9

This example illustrates an alternative procedure for the preparation of ureas of the present invention, exemplified by N-methyl-N-(4-morpholinecarbonyl)-4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]aniline (9).




25

To a stirred solution of N-methyl-4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]aniline (50 mg, 0.183 mmol) and diisopropylethylamine (73.3 mL, 0.421 mmol) in dichloromethane (2 mL) was added 4-morpholinecarbonyl chloride (21 mL, 0.183 mmol). Stirring was continued for 100 hours at room temperature. At the completion of the reaction, the reaction mixture was diluted with ethyl acetate, and washed sequentially with 1N HCl and brine. The organic layer was dried over MgSO₄ and filtered. Solvents were removed from the filtrate under reduced pressure and the residue was purified by flash chromatography on silica gel (hexanes and ethyl acetate (4:1) as eluant) to provide the title compound (50 mg, 70.0%) as a clear oil.


¹H (CDCl₃): δ 7.69 (d, *J*=8.4 Hz, 2H), 7.12 (d, *J*=8.4 Hz, 2H), 3.95 (bs, 1H), 3.50 (m, 4H), 3.25 (s, 3H), 3.20 (m, 4H). MS (ES⁻): 387 (M+H, 100).

EXAMPLE 10

This example provides procedures for the preparation of N-alkyl derivatives of the sulfonamide compounds of the invention provided in Example 2, and related analogs.



Synthesis of compound 10.1




10.1

To a stirred mixture of sulfonamide **2** (2.08 g, 5.21 mmol, from Example 2) and ethyl bromoacetate (0.950 g, 5.68 mmol) in DMF (15 mL) and THF (5 mL), was added sodium hydride (60% dispersion in mineral oil, 0.230 g, 5.70 mmol) at 0°C. The mixture was stirred overnight as the temperature was allowed to warm to room temperature. At the completion of the reaction (as determined by thin-layer chromatography), the mixture was diluted with ethyl acetate, washed with brine (3X), dried over MgSO₄, and concentrated. The residue was purified by flash chromatography on silica gel (hexane/EtOAc gradient from 8:1 to 2:1, as eluant) to give the title compound (2.01 g, 84%) as a clear oil.

¹H NMR (CDCl₃): δ 7.69 (d, *J*=7.6 Hz, 2H), 7.66 (d, *J*=8.4 Hz, 2H), 7.60 (t, *J*=7.6 Hz, 1H), 7.48 (t, *J*=7.6 Hz, 2H), 7.32 (d, *J*=8.4 Hz, 2H), 4.44 (s, 2H), 4.15 (q, *J*=6.5 Hz, 2H), 3.75 (s, 1H), 1.21 (t, *J*=6.5 Hz, 3H). MS (ES⁻): 484 (M-H, 100). Anal. Calcd. for C₁₉H₁₇F₆NO₅S: C, 47.01; H, 3.53; N, 2.89; S, 6.61. Found: C, 47.10; H, 3.59; N, 2.81; S, 6.64.

Synthesis of compound 10.2




10.2

To a solution of compound **10.1** (100 mg, 0.176 mmol) in 1 mL methanol, 1 mL THF, and 1 mL water was added 1 N NaOH (0.440 mL). The resulting mixture was stirred at room temperature for 3 hours, then diluted into ethyl acetate, washed with brine (3X), dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by flash chromatography on SiO₂ (CH₂Cl₂:MeOH:AcOH (10:1:0.05) as eluant) to give the title compound (60 mg, 65% yield) as a clear oil.

¹H NMR (CD₃OD): δ 7.80 (m, 5H), 7.66 (t, *J*=7.2 Hz, 2H), 7.46 (d, *J*=8.2 Hz, 2H), 4.60 (s, 2H). MS (ES⁻): 456 (M-H, 100). Anal. Calcd. for C₁₇H₁₃F₆NO₅S: C, 44.65; H, 2.87; N, 3.06; S, 7.01. Found: C, 44.84; H, 2.96; N, 3.21.

Synthesis of compound 10.3




10.3

A portion of the compound from Example 10.1 (71 mg, 0.146 mmol) was treated with LiAlH_4 (1 M in THF, 200 mL, 0.200 mmol) in THF at 0°C . The reaction was quenched by careful addition of water. Ethyl acetate was added and the resulting suspension was filtered through a Celite pad. The filtrate was washed with brine, dried over MgSO_4 , and concentrated under reduced pressure. Crude product was purified by chromatography on SiO_2 (hexanes/EtOAc (2:1) as eluant) to provide the title compound (48 mg, 74% yield).

^1H NMR (CDCl_3): δ 7.68 (d, $J=8.6$ Hz, 2H), 7.62 (d, $J=8.0$ Hz, 2H), 7.60 (t, $J=7.2$ Hz, 1H), 7.50 (dd, $J=8.0, 7.2$ Hz, 2H), 7.20 (d, $J=8.6$ Hz, 2H), 3.73 (m, 4H), 3.60 (s, 1H). MS (ES $^-$): 442 (M-H, 100). Anal. Calcd. for $\text{C}_{17}\text{H}_{15}\text{F}_6\text{NO}_4\text{S}$: C, 46.05; H, 3.41; N, 3.16; S, 7.23. Found: C, 46.15; H, 3.44; N, 3.21.

Synthesis of compound 10.4



10.4

A solution of the acid (10.2) from above (137 mg, 0.300 mmol) in 4 mL of CH_2Cl_2 with two drops of DMF added was treated with a solution of oxalyl chloride (2M in CH_2Cl_2 , 4 mL) at room temperature for 2 hours. Solvent was evaporated under reduced pressure to provide an intermediate acyl chloride. To a solution of acyl chloride in CH_2Cl_2 was added pyrrolidine (80 (L, 0.958 mmol). After stirring for 30 min, additional CH_2Cl_2 was added. The organic layer was washed with 1 N HCl (2X) and brine (3X), dried over MgSO_4 , and concentrated. The crude

product was purified by flushing chromatography on SiO₂ (4:1 hexanes:EtOAc as eluant) to provide 150 mg of the title compound (100%).

¹H NMR (CDCl₃): δ 7.68 (d, *J*=8.6 Hz, 2H), 7.60 (d, *J*=7.6 Hz, 2H), 7.55 (t, *J*=7.6 Hz, 1H), 7.42 (t, *J*=7.6 Hz, 2H), 7.35 (t, *J*=7.6 Hz, 2H), 4.45 (s, 2H), 4.40 (s, 1H), 3.52 (t, *J*=5.5 Hz, 2H), 3.38 (t, *J*=5.5 Hz, 2H), 1.95 (m, 2H), 1.82 (m, 2H). MS (ES⁻): 509 (M-H, 100). Anal. Calcd. for C₂₁H₂₀F₆N₂O₄S: C, 49.41; H, 3.95; N, 5.49; S, 6.28. Found: C, 49.59; H, 4.00; N, 5.42; S, 6.25.

The following compounds were prepared in accordance with the above procedure, replacing pyrrolidine with the appropriate amine:

10.5 (using diethylamine)

¹H NMR (CDCl₃): δ 7.65 (d, *J*=8.5 Hz, 2H), 7.60 (d, *J*=7.5 Hz, 2H), 7.55 (t, *J*=7.5 Hz, 1H), 7.45 (d, *J*=7.5 Hz, 2H), 7.31 (d, *J*=8.5 Hz, 2H), 4.38 (s, 2H), 3.97 (s, 1H), 3.42 (q, *J*=6.6 Hz, 2H), 3.25 (q, *J*=6.6 Hz, 2H), 1.21 (t, *J*=6.6 Hz, 3H), 0.90 (t, *J*=6.6 Hz, 3H). MS (ES⁺): 513 (M+H, 100). Anal. Calcd. for C₂₁H₂₂F₆N₂O₄S: C, 49.22; H, 4.33; N, 5.47; S, 6.26. Found: C, 49.48; H, 4.35; N, 5.37; S, 6.24.

10.6 (using 4-methylpiperazine)

¹H NMR (CDCl₃): δ 7.62 (d, *J*=8.5 Hz, 2H), 7.60 (t, *J*=7.5 Hz, 1H), 7.55 (t, *J*=7.5 Hz, 2H), 7.45 (t, *J*=7.5 Hz, 2H), 7.25 (d, *J*=7.5 Hz, 2H), 4.45 (s, 2H), 3.70 (m, 2H), 3.45 (m, 2H), 2.50 (m, 2H), 2.35 (m, 2H), 2.30 (s, 3H). MS (ES⁻): 540 (M-H, 100), MS (ES⁻): 538 (M-H, 100). Anal. Calcd. for C₂₂H₂₃F₆N₃O₄S: C, 48.98; H, 4.30; F, 21.13; N, 7.79; O, 11.86; S, 5.94. Found: C, 49.26; H, 4.38; N, 7.64; S, 5.83.


10.7 (using 2-methylpiperidine)

¹H NMR (CDCl₃): δ 7.62 (d, *J*=8.5 Hz, 2H), 7.60 (t, *J*=7.5 Hz, 1H), 7.60 (d, *J*=7.5 Hz, 2H), 7.45 (t, *J*=7.5 Hz, 2H), 7.30 (d, *J*=8.5 Hz, 2H), 4.50 (s, 2H), 4.30 (m, 1H), 3.85 (m, 1H), 1.40-1.70 (m, 6H), 0.80 (d, *J*=5.5 Hz, 3H). MS (ES⁻): 537 (M-H, 100). Anal. Calcd. for C₂₃H₂₄F₆N₂O₄S: C, 51.30; H, 4.49; N, 5.20; S, 5.95. Found: C, 51.81; H, 4.68; N, 5.13; S, 5.89.

EXAMPLE 11

This example illustrates a procedure for the reduction of the amides prepared in Example 10 to their corresponding amine derivatives.

Synthesis of compound 11.1 from compound 10.4:



The amide produced in 10.4 (126 mg, 0.247 mmol) was treated with
 5 LiAlH_4 (18.7 mmol, 37.4 mmol) in THF (3 mL) at reflux for 2 hours. The reaction
 was quenched by careful addition of water, ethyl acetate was added, and the resulting
 suspension was filtered through a Celite pad. The filtrate was washed with brine,
 dried over MgSO_4 , and stripped. Crude product was purified by flushing
 chromatography on SiO_2 eluted CH_2Cl_2 :MeOH (20:1) to give 60 mg (50%) of amine
 10 11.1.

^1H NMR (CDCl_3): δ 7.60 (t, $J=7.6$ Hz, 1H), 7.57 (d, $J=8.5$ Hz, 2H), 7.52
 (d, $J=7.6$ Hz, 2H), 7.45 (t, $J=7.6$ Hz, 2H), 7.11 (d, $J=8.5$ Hz, 2H), 3.80 (t, $J=6.5$ Hz,
 2H), 2.70 (m, 2H), 2.55 (m, 4H), 1.80 (m, 4H). MS (ES+): 497 (M+H, 100). Anal.
 15 Calcd. for : C, 50.80; H, 4.47; N, 5.64; S, 6.46; Found: C, 51.06; H, 4.58; N, 5.51; S,
 6.42.

In a similar manner, the amides 10.5, 10.6 and 10.7 were reduced to
 their corresponding amines.

20 Compound 11.2 (prepared from compound 10.5)

^1H NMR (CDCl_3): δ 7.55 (m, 5H), 7.45 (t, $J=7.5$ Hz, 2H), 7.15 (t, $J=8.5$
 Hz, 2H), 3.70 (m, 2H), 2.40-2.65 (m, 6H), 0.95 (s, 3H). MS (ES+): 499 (M+H, 100).
 Anal. Calcd. for $\text{C}_{21}\text{H}_{24}\text{F}_6\text{N}_2\text{O}_3\text{S}$: C, 50.60; H, 4.85; N, 5.62; S, 6.43. Found: C, 50.65;
 H, 4.90; N, 5.59; S, 6.35.

25

Compound 11.3 (prepared from compound 10.6)


^1H NMR (CDCl_3): δ 7.65 (d, $J=8.5$ Hz, 2H), 7.60 (m, 3H), 7.45 (t,
 $J=7.5$ Hz, 2H), 7.12 (d, $J=8.5$ Hz, 2H), 3.70 (t, $J=6.3$ Hz, 2H), 2.30-2.50 (m, 8H), 2.20
 (s, 3H). MS (ES-): 524 (M-H, 100). Anal. Calcd. for $\text{C}_{22}\text{H}_{25}\text{F}_6\text{N}_3\text{O}_3\text{S}$: C, 50.28; H,
 30 4.80; N, 8.00; S, 6.10. Found: C, 49.28; H, 5.12; N, 7.59; S, 5.74.

Compound 11.4 (prepared from compound 10.7)

¹H NMR (CDCl₃): δ 7.65 (d, *J*=8.6 Hz, 2H), 7.57 (d, *J*=7.4 Hz, 2H), 7.56 (t, *J*=7.4 Hz, 1H), 7.46 (t, *J*=7.4 Hz, 2H), 7.17 (d, *J*=8.6 Hz, 2H), 3.65 (m, 2H), 2.76 (m, 1H), 2.45 (m, 1H), 2.25 (m, 1H), 2.15 (m, 1H), 1.20-1.70 (m, 6H), 0.92 (d, *J*=5.0 Hz, 3H). MS (ES⁺): 525 (M+H, 100). Anal. Calcd. for C₂₃H₂₆F₆N₂O₃S: C, 52.67; H, 5.00; N, 5.34; S, 6.11. Found: C, 52.96; H, 5.10; N, 5.18; S, 5.89.


EXAMPLE 12

This example illustrates the synthesis of compound 12.



12

12.1 Preparation of *N*-trifluoroethylaniline derivative



To a suspension of 4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]aniline (9.07 g, 35.0 mmol) in CH₂Cl₂ (100 mL) was added a solution of trifluoroacetic anhydride (5.7 mL, 40.2 mmol) in CH₂Cl₂ (50 mL) dropwise at room temperature. The solution was stirred for 3 hours (solution cleared and TLC indicated that the reaction was completed). The reaction mixture was washed with water, aqueous NaHCO₃, and brine. The organic layer was drawn off, dried over MgSO₄, filtered, and concentrated to give 12.1 g of the intermediate trifluoroacetanilide (12.1a). The intermediate 12.1a was taken up in THF (50 mL) and treated with LiAlH₄ (4.00 g, 106 mmol) at reflux for 10 hours. The reaction was quenched by sequentially adding 4 mL water, 4 mL of 15% NaOH, and 12 mL of water. The resulting suspension was stirred for an additional 30 minutes, filtered through a Celite pad, which was then rinsed with THF. The combined filtrate and

rinse was concentrated under reduced pressure. The residue was taken up in EtOAc, washed with brine, dried over MgSO_4 , filtered, and concentrated. The resulting crude product was purified by chromatography on SiO_2 (4:1 hexanes:EtOAc as eluant) to provide 11.0 g (92%) of the title compound (**12.1b**).


¹H NMR (CDCl_3): δ 7.52 (d, $J=8.6$ Hz, 2H), 6.72 (d, $J=8.6$ Hz, 2H), 4.10 (bs, 1H), 3.80 (q, $J=8.5$ Hz, 2H), 3.31 (bs, 1H). MS (ES⁺): 342 (M+H, 100).

12.2 Sulfonylation of **12.1b**

A sample of **12.1b** from above (1.87 g, 5.48 mmol) was treated with benzenesulfonyl chloride (1.18 g, 6.68 mmol) in pyridine (10 mL) at room temperature for 10 days. The reaction mixture was diluted with EtOAc, washed with aqueous NaHCO_3 , and brine. The organic layer was dried over MgSO_4 , filtered, and concentrated. The crude product was purified by chromatography on SiO_2 (4:1 hexanes:EtOAc as eluant) to provide 1.65 g (62%) of compound **12**.

¹H NMR (CDCl_3): δ 7.78 (d, $J=8.8$ Hz, 2H), 7.61 (t, $J=7.6$ Hz, 1H), 7.58 (d, $J=7.6$ Hz, 2H), 7.46 (t, $J=7.6$ Hz, 2H), 4.24 (q, $J=8.2$ Hz, 2H), 3.41 (s, 1H). MS (ES⁻): 480 (M-H, 100). Anal. Calcd. for $\text{C}_{17}\text{H}_{12}\text{F}_9\text{NO}_3\text{S}$: C, 42.42; H, 2.51; N, 2.91; S, 6.66. Found: C, 42.70; H, 2.55; N, 2.84; S, 6.61.

In a similar manner, compound **12.3** was prepared by substituting 3-cyanobenzenesulfonyl chloride for benzenesulfonyl chloride.




12.3

¹H NMR (CDCl_3): δ 7.91 (s, 1H), 7.90 (d, $J=7.5$ Hz, 2H), 7.80 (d, $J=7.5$ Hz, 1H), 7.75 (d, $J=8.8$ Hz, 2H), 7.80 (d, $J=7.5$ Hz, 1H), 7.75 (d, $J=8.8$ Hz, 2H), 7.62 (t, $J=7.5$ Hz, 1H), 7.19 (d, $J=8.8$ Hz, 2H), 4.30 (q, $J=8.2$ Hz, 2H), 3.62 (s, 1H). MS (ES⁻): 505 (M-H, 100). Anal. Calcd. for $\text{C}_{18}\text{H}_{11}\text{F}_9\text{N}_2\text{O}_3\text{S}$: C, 42.70; H, 2.19; N, 5.53; S, 6.33. Found: C, 43.41; H, 2.46; N, 5.67.

EXAMPLE 13

This example illustrates the preparation of 13, and related derivatives 13.3 and 13.4.



Synthesis of compound 13.1

To a stirred mixture of indoline (5.32 g, 44.6 mmol) and toluenesulfonic acid (75 mg, 0.395 mmol) was added hexafluoroacetone hydrate (7 mL, 49.1 mmol) at 90°C over 15 min. The resulting mixture was heated at 135 °C overnight. After cooling to room temperature, the mixture was diluted with EtOAc, washed with aqueous NaHCO₃ (2X) and brine (2X). The organic layer was dried over MgSO₄, filtered, and concentrated. The crude product was purified by chromatography on SiO₂ (6:1 to 4:1 hexanes:EtOAc as eluant) to give the intermediate substituted indoline (13.1, 6.35 g, 50% yield).

¹H NMR (CDCl₃): δ 7.41 (s, 1H), 7.32 (d, *J*=8.5 Hz, 1H), 6.65 (d, *J*=8.5 Hz, 1H), 3.60 (d, *J*=5.5 Hz, 2H), 3.08 (d, *J*=5.5 Hz, 2H). MS (ES⁺): 286 (M+H, 100).


Synthesis of compound 13.

Sulfonylation of compound 13.1 was carried out as described in the Examples above to provide the title compound (13).

¹H NMR (CDCl₃): δ 7.84 (d, *J*=8.0 Hz, 2H), 7.68 (d, *J*=8.4 Hz, 2H), 7.59 (t, *J*=8.0 Hz, 1H), 7.52 (d, *J*=8.4 Hz, 1H), 7.48 (t, *J*=8.0 Hz, 2H), 7.41 (s, 1H), 3.97 (t, *J*=5.6 Hz, 2H), 3.38 (s, 1H), 3.00 (t, *J*=5.6 Hz, 2H). MS (ES⁻): 424 (M-H, 100). Anal. Calcd. for C₁₇H₁₃F₆NO₃S: C, 48.00; H, 3.08; N, 3.29. Found: C, 48.17; H, 3.06; N, 3.37.

In a similar manner, the following compounds were prepared, beginning with the tetrahydroquinoline and the appropriate (substituted)benzenesulfonyl chlorides.

13.3




13.3

5

^1H NMR (CDCl_3): δ 7.86 (d, $J=8.5$ Hz, 1H), 7.62 (d, $J=7.2$ Hz, 2H), 7.55 (t, $J=7.5$ Hz, 1H), 7.48 (d, $J=8.5$ Hz, 1H), 7.41 (t, $J=7.2$ Hz, 2H), 7.36 (s, 1H), 3.81 (dd, $J=6.5, 4.5$ Hz, 2H), 3.40 (s, 1H), 2.52 (dd, $J=7.0, 4.5$ Hz, 2H), 1.70 (m, 2H). MS (ES $^-$): 438 (M-H, 100). Anal. Calcd. for $\text{C}_{18}\text{H}_{15}\text{F}_6\text{NO}_3\text{S}$: C, 49.20; H, 3.44; N, 3.19; S, 7.30. Found: C, 49.55; H, 3.36; N, 3.31; S, 7.53.

10

13.4



13.4


15

^1H NMR (CDCl_3): δ 7.90 (s, 1H), 7.81 (d, $J=8.5$ Hz, 2H), 7.80 (d, $J=7.5$ Hz, 1H), 7.55 (d, $J=7.5$ Hz, 1H), 7.50 (t, $J=7.5$ Hz, 1H), 7.36 (s, 1H), 3.82 (dd, $J=6.5, 5.5$ Hz, 2H), 3.35 (s, 1H), 2.55 (t, $J=5.5$ Hz, 2H), 1.72 (m, 2H). MS (ES $^-$): 577 (M-H, 100). Anal. Calcd. for $\text{C}_{19}\text{H}_{14}\text{F}_6\text{N}_2\text{O}_3\text{S}$: C, 49.14; H, 3.04; N, 6.03; S, 6.91. Found: C, 49.25; H, 3.08; N, 5.86; 6.84.

20

EXAMPLE 14

This example illustrates the synthesis of compound 14.1.



5


Hexafluoroacetone trihydrate (9.68 g, 44 mmol) was added to 2,6-dimethylaniline (4.84 g, 40 mmol) and p-toluenesulfonic acid (304 mg, 1.6 mmol) at room temperature. The mixture was stirred at 130°C for 14 hours under a nitrogen atmosphere. After cooling down, the solidified mixture was washed with sodium bicarbonate solution, water, and hexane and dried to afford 10g of 4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]-2,6-dimethylaniline. ¹H NMR (DMSO): δ 8.11 (s, 1H), 7.08 (s, 2H), 4.89 (s, 2H), 2.10 (s, 6H). MS (ES) *m/z* 288.1 ([M+H]⁺).

To a solution of 4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]-2,6-dimethylaniline (4.3 g, 15 mmol) and 2,6-lutidine (3.5 mL, 30 mmol) in acetone (25 mL), was added benzenesulfonyl chloride (3.18 g, 18 mmol). The mixture was stirred at room temperature for 20 hours, at 50°C for 20 hours, and at 70°C for 24 hours. After cooling down, the solid was filtered off. The filtrate was concentrated, and the residue was treated by 3N HCl (200 mL), and it was extracted with ethyl acetate. The organic layer was separated, washed with saturated sodium bicarbonate solution and brine, dried, and concentrated. The residue was purified by silica column chromatography (eluted by 9:1 DCM/EtOAc) to afford 5.1 g of product (example 14.1). ¹H NMR (DMSO): δ 9.52 (s, 1H), 8.64 (s, 1H), 7.69 (m, 3H), 7.59 (m, 2H), 7.31 (s, 2H), 1.99 (s, 6H). MS (ES) *m/z* 426.0 ([M-H]⁺). Anal. Calcd. for C₁₇H₁₅F₆NO₃S: C, 47.78; H, 3.54; N, 3.28. Found: C, 47.69; H, 3.45; N, 3.33.

25

Compounds 14.2-14.11 were prepared from their corresponding anilines in a manner similar to that described for 14.1.

14.2




14.2

5

^1H NMR (DMSO): δ 9.75 (s, 1H), 8.42 (s, 1H), 7.72 (m, 2H), 7.66 (m, 1H), 7.57 (m, 2H), 7.18 (s, 1H), 6.91 (s, 1H), 2.40 (s, 3H), 1.97 (s, 3H). MS (EI) m/z 427 (M^+). Anal. Calcd. for $\text{C}_{17}\text{H}_{15}\text{F}_6\text{NO}_3\text{S}$: C, 47.78; H, 3.54; N, 3.28. Found: C, 47.60; H, 3.57; N, 3.33.

10

14.3




14.3

^1H NMR (DMSO): δ 9.59 (s, 1H), 8.65 (s, 1H), 7.72 (m, 2H), 7.66 (m, 1H), 7.58 (m, 2H), 7.39 (s, 1H), 7.29 (s, 1H), 3.17 (h, $J=6.9$ Hz, 1H), 1.98 (s, 3H), 0.93 (d, $J=6.9$ Hz, 6H). MS (EI) m/z 455 (M^+). Anal. Calcd. for $\text{C}_{19}\text{H}_{19}\text{F}_6\text{NO}_3\text{S}$: C, 50.11; H, 4.21; N, 3.08. Found: C, 50.18; H, 4.23; N, 3.06.

15

14.4




14.4

20

^1H NMR (CDCl_3): δ 7.78 (m, 3H), 7.65-7.50 (m, 4H), 7.34 (d, $J=8.5$ Hz, 1H), 3.54 (s, 1H), 3.23 (s, 3H). MS (EI) m/z 447 (M^+). Anal. Calcd. for $\text{C}_{16}\text{H}_{12}\text{ClF}_6\text{NO}_3\text{S}$: C, 42.92; H, 2.70; N, 3.13; S, 7.16. Found: C, 43.19; H, 2.80; N, 3.16; S, 7.27.

25

14.5




14.5

5 $^1\text{H-NMR}$ (DMSO): δ 9.80 (s, 1H), 8.68 (s, 1H), 7.78 (d, $J=7.12, 1.56$ Hz, 2H), 7.56 (m, 3H), 7.33 (d, $J=8.36, 1\text{H}$), 7.15 (d, $J=12.00$ Hz, 2H), 3.59 (s, 3H). MS (ES⁺): 430 (M+H, 75). Anal. Calc. for $\text{C}_{16}\text{H}_{13}\text{F}_6\text{NO}_4\text{S}$: C, 44.76; H, 3.05; N, 3.26; S, 7.47. Found: C, 44.90; H, 3.10; N, 3.32; S, 7.57.

10

14.6




14.6

$^1\text{H-NMR}$ (DMSO): δ 9.34 (s, 1H), 8.71 (s, 1H), 7.60(m, 5H), 7.12 (s, 1H), 6.91 (s, 1H), 3.17 (s, 3H), 2.33 (s, 3H). MS (ES⁺): 444 (M+H, 17). MS (ES⁻): 442 (M-H, 100). Anal. Calc. for $\text{C}_{17}\text{H}_{15}\text{F}_6\text{NO}_4\text{S}$: C, 46.05; H, 3.41; N, 3.16; S, 7.23. Found: C, 46.18; H, 3.50; N, 3.16; S, 7.33.

15

14.7




14.7

20

$^1\text{H-NMR}$ (DMSO): δ 10.22 (s, 1H), 8.69 (s, 1H), 7.50 (m, 2H), 7.20 (m, 2H), 2.21 (s, 3H). MS (ES⁺): 488 (M+H, 15). Anal. Calc. for $\text{C}_{14}\text{H}_9\text{Cl}_2\text{F}_6\text{NO}_3\text{S}_2$:

C, 34.44; H, 1.86; N, 2.87; Cl, 4.52; S, 13.13. Found: C, 34.65; H, 1.80; N, 2.69; Cl, 14.44; S, 13.02.


14.8



14.8

¹H-NMR (DMSO): δ 9.90 (s, 1H), 8.65 (s, 1H), 7.6 (m, 6H), 7.36 (d, J=8.40 Hz, 1H), 7.08 (d, J=8.48 Hz, 1H), 3.17 (m, 1H), 0.90 (d, J=6.80 Hz, 6H). MS (ES⁻): 440 (M-H, 100). Anal. Calc. for C₁₈H₁₇F₆NO₃S: C, 48.98; H, 3.88; N, 3.17; S, 7.26. Found: C, 49.03; H, 3.83; N, 3.18; S, 7.37.


14.9



14.9

¹H-NMR (DMSO): δ 10.14 (s, 1H), 8.69 (s, 1H), 8.15 (d, J=7.76 Hz, 1H), 8.01 (m, 2H), 7.81 (t, J=7.72 Hz, 1H), 7.54 (s, 1H), 7.40 (d, J=8.44 Hz, 1H), 7.04 (d, J=8.64 Hz, 1H), 3.15 (m, 1H), 0.94 (d, J=6.88 Hz, 6H). MS (ES⁻): 465 (M-H, 100). Anal. Calc. for C₁₉H₁₆F₆N₂O₃S: C, 48.93; H, 3.46; N, 6.01; S, 6.88. Found: C, 48.99; H, 3.39; N, 5.84; S, 6.82.

14.10




14.10

¹H-NMR (DMSO): δ 9.44 (s, 1H), 8.71 (s, 1H), 7.90 (m, 2H), 7.70 (m, 4H), 7.30 (m, 1H), 6.65 (d, J=8.44 Hz, 1H), 1.43 (s, 9H). MS (ES⁻): 454 (M-H, 100).

Anal. Calc. for C₁₉H₁₉F₆NO₃S: C, 50.11; H, 4.21; N, 3.08; S, 7.04. Found: C, 50.18;

5 H, 4.22; N, 3.07; S, 7.04.

14.11




14.11

10 ¹H-NMR (DMSO): δ 11.06 (s, 1H), 8.64 (s, 1H), 7.60 (d, J=9Hz, 2H), 7.39 (s, 1H), 7.25 (d, J=9Hz, 2H). Anal. Calc. for C₁₄H₉Cl₂F₆NO₃S₂: C, 32.92; H, 1.49; N, 2.95; S, 13.52. Found: C, 33.06; H, 1.44; N, 2.89; S, 13.53.

15

EXAMPLE 15

This example illustrates the preparation of compound 15.1



15.1

20 Fuming nitric acid (90%, 5 mL) was added drop wise to 1,1,1,3,3,3-hexafluoro-2-phenyl-2-propanol (4.2 mL, 25 mmol) in concentrated sulfuric acid (15 mL) at 0°C. The mixture was stirred at 0°C for 30 minutes, and at room temperature for 2 hours. Then it was poured into ice water. The solid was filtered, washed with sodium bicarbonate solution, water, and hexane, and dried to give 6.2 g of 1-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]-3-nitrobenzene. ¹H NMR (CDCl₃): δ


25

8.66 (s, 1H), 8.38 (d, J=8 Hz, 1H), 8.06 (d, J=8 Hz, 1H), 7.69 (t, J=8 Hz, 1H), 3.66 (bs, 1H). MS (EI) m/z 289 (M^+).

Palladium on carbon (10%, 150 mg) was added to the product above (2.0 g, 6.9 mmol) in ethanol (20 mL). The mixture was stirred under 1 atmosphere of hydrogen at room temperature for 6 hours. The catalyst was filtered off through celite. The filtrate was concentrated to give 1.78 g of solid product (3-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]aniline). ^1H NMR (DMSO): δ 8.36 (s, 1H), 7.11 (t, J=8 Hz, 1H), 6.93 (s, 1H), 6.78 (d, J=8 Hz, 1H), 6.66 (d, J=8 Hz, 1H), 5.40 (bs, 1H). MS (EI) m/z 259 (M^+).

To a solution of the above product (770 mg, 3 mmol) and 2,6-lutidine (0.7 mL, 6 mmol) in acetone (10 mL), was added benzenesulfonyl chloride (0.46 mL, 3.6 mmol). The mixture was stirred at room temperature for 20 hours. The solid was filtered off. The filtrate was concentrated, and the residue was treated by 3N HCl (100 mL), and it was extracted with ethyl acetate. The organic layer was separated, washed with saturated sodium bicarbonate solution and brine, dried, and concentrated. The residue was purified by silica column chromatography (eluted by 1:2 EtOAc/Hexane) to afford 1.08 g of **15.1**. ^1H NMR (CDCl_3): δ 7.73 (d, J=7.2 Hz, 2H), 7.54 (m, 1H), 7.45 (m, 3H), 7.35 (m, 2H), 7.24 (s, 1H), 6.52 (s, 1H), 3.39 (s, 1H). MS (ES) m/z 398.0 ($[\text{M}-\text{H}]^-$). Anal. Calcd. for $\text{C}_{15}\text{H}_{11}\text{F}_6\text{NO}_3\text{S}$: C, 45.12; H, 2.78; N, 3.51; S, 8.03. Found: C, 45.27; H, 2.88; N, 3.49; S, 8.05.

Compound **15.2** was prepared in a manner the same as that described above except that 3-(2,5-dimethylthienyl)sulfonyl chloride was used instead of benzenesulfonyl chloride.



15.2


^1H NMR (CDCl_3): δ 7.51 (m, 1H), 7.39 (m, 2H), 7.23 (m, 1H), 6.81 (s, 1H), 6.60 (s, 1H), 3.66 (s, 1H), 2.35 (s, 3H), 2.33 (s, 3H). MS (EI) m/z 433 (M^+).

Anal. Calcd. for $C_{15}H_{13}F_6NO_3S_2$: C, 41.57; H, 3.02; N, 3.23. Found: C, 41.70; H, 3.02; N, 3.20.

5

EXAMPLE 16

This example illustrates the preparation of compound **16.1**

**16.1**


10 Sodium hydride (60 %, 116.4 mg, 2.91 mmol) was added to compound **14.1** (1.13 g, 2.65 mmol) in DMF (10 mL) at room temperature. The mixture was stirred at room temperature for 20 minutes. Then iodomethane (0.18 ml, 2.91 mmol) was added. The mixture was stirred at room temperature for 2 hours. DMF was evaporated under vacuum, and the residue was treated with water, which was
15 extracted with EtOAc. The organic layer was separated, washed with brine, dried, and concentrated. The residue was purified by silica column chromatography (eluted by 1;5 EtOAc/Hexane) to afford 1.06 g of product.

1H NMR ($CDCl_3$): δ 7.82 (m, 2H), 7.60 (m, 1H), 7.52 (m, 2H), 7.38 (s, 2H), 3.85 (bs, 1H), 3.18 (s, 3H), 2.12 (s, 6H). MS (ES) m/z 440.0 ($[M-H]^-$). Anal.
20 Calcd. for $C_{18}H_{17}F_6NO_3S$: C, 48.98; H, 3.88; N, 3.17. Found: C, 49.12; H, 3.86; N, 3.26.

Compounds **16.2-16.38** were prepared in a manner similar to that described for **16.1**.

25

16.2




16.2

¹H NMR (DMSO): δ 8.55 (s, 1H), 7.77 (m, 1H), 7.68 (m, 4H), 7.38 (s, 1H), 6.53 (s, 1H), 3.07 (s, 3H), 2.36 (s, 3H), 2.26 (s, 3H). MS (EI) *m/z* 441 (M⁺).

Anal. Calcd. for C₁₈H₁₇F₆NO₃S: C, 48.98; H, 3.88; N, 3.17. Found: C, 49.20; H, 3.93; N, 3.16.


16.3



16.3


¹H NMR (DMSO): δ 8.73 (s, 1H), 7.85 (m, 2H), 7.74 (m, 1H), 7.66 (m, 2H), 7.45 (s, 1H), 7.37 (s, 1H), 3.14 (s, 3H), 2.90 (h, J=6.9 Hz, 1H), 2.05 (s, 3H), 1.06 (dd, J=6.9 Hz, 6H). MS (ES) *m/z* 468.1 ([M-H]⁻). Anal. Calcd. for C₂₀H₂₁F₆NO₃S: C, 51.17; H, 4.51; N, 2.98. Found: C, 51.18; H, 4.54; N, 3.10.

16.4



¹H NMR (CDCl₃): δ 7.65-7.30 (m, 9H), 3.40 (s, 1H), 3.19 (s, 3H). MS (ES) *m/z* 412.0 ([M-H]⁻). Anal. Calcd. for C₁₆H₁₃F₆NO₃S: C, 46.49; H, 3.17; N, 3.39; S, 7.76. Found: C, 46.74; H, 3.30; N, 3.35; S, 7.79.

16.5




16.5

¹H NMR (CDCl₃): δ 7.63 (m, 1H), 7.43 (m, 3H), 6.67 (s, 1H), 3.37 (s, 1H), 3.24 (s, 3H), 2.36 (s, 3H), 2.05 (s, 3H). MS (EI) *m/z* 447 (M⁺). Anal. Calcd. for C₁₆H₁₅F₆NO₃S₂: C, 42.95; H, 3.38; N, 3.13; S, 14.33. Found: C, 43.00; H, 3.29; N, 3.08; S, 14.44.

10

16.6




16.6

¹H-NMR (DMSO): δ 8.84 (s, 1H), 7.60 (m, 5H), 7.32 (d, J=8.92 Hz, 1H), 7.22(m, 2H), 3.44 (s, 1H), 3.12 (s, 1H). MS (ES⁻): 442 M-H, 100). Anal. Calc. for C₁₇H₁₅F₆NO₄S: C, 46.05; H, 3.41; N, 3.16; S, 7.23. Found: C, 46.14; H, 3.43; N, 3.15; S, 7.34.

15

16.7




16.7

20

¹H-NMR (DMSO): δ 8.78 (s, 1H), 7.70 (m, 3H), 7.60 (m, 2H), 7.17 (s, 1H), 7.01(s, 1H), 3.26 (s, 3H), 3.07 (s, 3H), 2.38 (s, 3H). MS (ES⁻): 456 (M-H, 100). Anal. Calc. for C₁₈H₁₇F₆NO₄S: C, 47.27; H, 3.75; N, 3.06; S, 7.01. Found: C, 47.32; H, 3.75; N, 3.07; S, 7.08.

25

16.8




16.8

5 $^1\text{H-NMR}$ (DMSO): δ 8.80 (s, 1H), 7.63 (s, 1H), 7.50 (d, $J=8.52$ Hz, 1H), 7.32 (s, 1H), 7.14 (d, $J=8.52$ Hz, 1H), 3.23 (s, 3H), 2.37 (s, 3H). MS (ES⁺): 502 (M+H, 100). Anal. Calc. for $\text{C}_{15}\text{H}_{11}\text{Cl}_2\text{F}_6\text{NO}_3\text{S}_2$: C, 35.87; H, 2.21; N, 2.79; Cl, 14.12; S, 12.77. Found: C, 35.95; H, 2.10; N, 2.63; Cl, 14.12; S, 12.88.

10


16.9



16.9

$^1\text{H-NMR}$ (DMSO): δ 8.77 (s, 1H), 7.62 (d, $J=8.44$ Hz, 2H), 7.50 (m, 1H), 7.40 (d, $J=8.96$ Hz, 2H), 7.36 (s, 1H), 7.31 (s, 1H), 7.0 (m, 1H), 4.96 (m, 2H), MS
15 (ES⁻): 568 (M-H, 56). Anal. Calc. for $\text{C}_{18}\text{H}_{11}\text{Cl}_2\text{F}_6\text{NO}_3\text{S}_3$: C, 37.90; H, 1.94; N, 2.46; Cl, 12.43; S, 16.87. Found: C, 37.84; H, 1.81; N, 2.38; Cl, 12.52; S, 16.96.

16.10




16.10

20

¹H-NMR (DMSO): δ 8.74 (s, 1H), 8.36 (d, J=8.4 Hz, 1H), 7.94 (d, J=8.04 Hz, 1H), 7.88 (d, J=8.4 Hz, 1H), 7.70 (m, 1H), 7.6 (m, 6H), 7.42 (s, 1H), 5.29(s, 2H). MS (ES-): 613 (M-H, 83). Anal. Calc. for C₂₃H₁₄Cl₂F₆N₂O₃S₂: C, 44.89; H, 2.29; N, 4.55; Cl, 11.52; S, 10.42. Found: C, 45.32; H, 2.33; N, 4.51; Cl, 11.31; S, 10.22.

16.11




16.11

¹H-NMR (DMSO): δ 8.78 (s, 1H), 7.64 (d, J=8.8 Hz, 2H), 7.40(m, 4H), 6.9(m, 2H), 5.17 (s, 2H). MS (ES-): 568 (M-H, 20). Anal. Calc. for C₁₈H₁₁Cl₂F₆NO₃S₃: C, 37.90; H, 1.94; N, 2.46; Cl, 12.43; S, 16.87. Found: C, 37.63; H, 1.88; N, 2.45; Cl, 12.68; S, 16.77.

15


16.12



16.12

¹H-NMR (DMSO): δ 8.82 (s, 1H), 7.68 (d, J=8.44 Hz, 2H), 7.47(d, J=8.8 Hz, 2H), 7.34 (s, 1H), 6.8(m, 3H), 6.58(m, 1H), 4.25(m, 2H), 4.06(m, 3H), MS (ES-): 620 (M-H, 100). Anal. Calc. for C₂₂H₁₅Cl₂F₆NO₅S₂: C, 42.46; H, 2.43; N, 2.25; Cl, 11.39; S, 10.30. Found: C, 42.55; H, 2.34; N, 2.27; Cl, 11.46; S, 10.22.


16.13



16.13

¹H-NMR (DMSO): δ 8.77 (s, 1H), 7.62 (d, J=8.44 Hz, 2H), 7.40 (m, 3H), 6.70(m, 3H), 4.85(s, 2H), 4.17(s, 4H). MS (ES⁻): 620 (M-H, 100). Anal. Calc. for C₂₂H₁₅Cl₂F₆NO₂S₂: C, 42.46; H, 2.43; N, 2.25; Cl, 11.39; S, 10.30. Found: C, 42.54; H, 2.47; N, 2.18; Cl, 11.35; S, 10.39.


16.14



16.14

¹H-NMR (DMSO): δ 8.73 (s, 1H), 7.58 (d, J=8.8 Hz, 2H), 7.28(m, 7H), 6.97 (d, J=1.12 Hz, 1H), 4.86 (s, 2H), 2.41 (s, 3H), 1.93 (s, 3H). MS (ES⁺): 524 (M+H, 100). Anal. Calc. for C₂₂H₁₉F₆NO₃S₂: C, 50.47; H, 3.66; N, 2.68; S, 12.25. Found: C, 50.57; H, 3.58; N, 2.60; S, 12.35.


16.15



16.15


¹H-NMR (DMSO): δ 8.84 (s, 1H), 7.44 (d, J=8.44 Hz, 2H), 7.24(d, J=8.68 Hz, 2H), 6.90 (s, 1H), 4.41 (m, 1H), 2.39 (s, 3H), 2.20 (s, 3H), 1.02 (d, J=6.72 Hz, 6H), MS (ES+): 476 (M+H, 100). Anal. Calc. for C₁₈H₁₉F₆NO₃S₂: C, 45.47; H, 4.03; N, 2.95; S, 13.49. Found: C, 45.66; H, 4.10; N, 2.90; S, 13.58.

5

16.16**16.16**

¹H-NMR (DMSO): δ 8.76 (s, 1H), 7.60 (d, J=8.52 Hz, 2H), 7.46(s, 1H), 7.40(m, 1H), 7.32(m, 3H), 7.24(t, 1H), 6.96 (s, 1H), 4.87 (s, 2H), 2.41 (s, 3H), 1.93 (s, 3H). MS (ES-): 600 (M-H, 60). Anal. Calc. for C₂₂H₁₈BrF₆NO₃S₂: C, 43.86; H, 3.01; N, 2.33; S, 10.65. Found: C, 43.83; H, 2.87; N, 2.36; S, 10.81.

10


16.17**16.17**

¹H-NMR (DMSO): δ 8.75 (s, 1H), 7.60 (d, J=8.45 Hz, 2H), 7.30 (d, J=8.88 Hz, 2H), 6.94(s, 1H), 6.73(m, 3H), 4.73(s, 2H), 4.16(s, 4H), 2.40 (s, 3H), 1.91 (s, 3H). MS (ES-): 580 (M-H, 80). Anal. Calc. for C₂₄H₂₁F₆NO₅S₂: C, 49.57; H, 3.64; N, 2.41; S, 11.03. Found: C, 49.39; H, 3.77; N, 2.48; S, 11.01.

15

20


16.18



16.18

¹H-NMR (DMSO): δ 8.79 (s, 1H), 7.70 (m, 6H), 7.40 (m, 1H), 6.76 (d, J=8.50 Hz, 1H), 3.49 (m, 1H), 3.11 (s, 3H), 1.20 (m, 6H). MS (ES⁻): 454 (M-H, 100).
 5 Anal. Calc. for C₁₉H₁₉F₆NO₃S: C, 50.11; H, 4.21; N, 3.08; S, 7.04. Found: C, 50.12; H, 4.18; N, 3.09; S, 7.12.


16.19



16.19

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.21 (m, 1H), 8.11 (s, 1H), 7.90 (d, J=8.08 Hz, 1H), 7.79 (m, 1H), 7.58 (m, 2H), 7.32 (m, 6H), 4.99 (s, 2H), MS (ES⁻):
 547 (M-H, 100). Anal. Calc. for C₂₃H₁₅ClF₆N₂O₃S: C, 50.33; H, 2.75; N, 5.10; Cl,
 10 6.46; S, 5.84. Found: C, 50.49; H, 2.80; N, 5.09; Cl, 6.54; S, 5.73.

16.20




16.20

¹H-NMR (DMSO): δ 8.73 (s, 1H), 8.21 (d, J=8.60 Hz, 1H), 8.07 (s, 1H), 7.81 (m, 2H), 7.54 (d, J=8.44 Hz, 2H), 7.26 (d, J=8.80 Hz, 2H), 7.10 (m, 3H),
 20 7.00 (m, 1H), 4.88 (s, 2H), 2.28 (s, 3H). MS (ES⁻): 527 (M-H, 90). Anal. Calc. for

$C_{24}H_{18}F_6N_2O_3S$: C, 54.55; H, 3.43; N, 5.30; S, 6.07. Found: C, 54.05; H, 3.46; N, 5.26; S, 5.97.


16.21



16.21

1H -NMR (DMSO): δ 8.74 (s, 1H), 8.21 (m, 1H), 8.10 (s, 1H), 7.87 (m, 2H), 7.58 (d, $J=8.48$ Hz, 2H), 7.28 (d, $J=8.84$ Hz, 2H), 7.13 (m, 1H), 7.02 (m, 3H), 4.85 (s, 2H), 2.21 (s, 3H). MS (ES⁻): 527 (M-H, 100). Anal. Calc. for $C_{24}H_{18}F_6N_2O_3S$: C, 54.55; H, 3.43; N, 5.30; S, 6.07. Found: C, 54.59; H, 3.47; N, 5.24; S, 6.00.


16.22



16.22

1H -NMR (DMSO): δ 8.77 (s, 1H), 8.30 (m, 1H), 8.15 (s, 1H), 7.84 (m, 2H), 7.60 (d, $J=8.44$ Hz, 2H), 7.27 (m, 6H), 4.92 (s, 2H). MS (ES⁻): 547 (M-H, 100). Anal. Calc. for $C_{23}H_{15}ClF_6N_2O_3S$: C, 50.33; H, 2.75; N, 5.10; Cl, 6.46; S, 5.84. Found: C, 50.26; H, 2.77; N, 5.07; Cl, 6.50; S, 5.82.


16.23



16.23

¹H-NMR (DMSO): δ 8.74 (s, 1H), 8.20 (d, J=7.32 Hz, 1H), 8.09 (s, 1H), 7.85 (m, 2H), 7.57 (d, J=8.12 Hz, 2H), 7.27 (d, J=8.40 Hz, 2H), 7.11 (m, 4H), 4.85 (s, 2H), 2.21 (s, 3H). MS (ES⁻): 527 (M-H, 60). Anal. Calc. for C₂₄H₁₈F₆N₂O₃S: C, 54.55; H, 3.43; N, 5.30; S, 6.07. Found: C, 54.56; H, 3.39; N, 5.29; S, 5.98.


16.24



16.24

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.21 (m, 1H), 8.13 (d, J=1.60 Hz, 1H), 7.83 (m, 2H), 4.92 (s, 2H), 7.58 (d, J=8.44 Hz, 2H), 7.30 (m, 6H), 4.90 (s, 2H). MS (ES⁻): 547 (M-H, 100). Anal. Calc. for C₂₃H₁₅ClF₆N₂O₃S: C, 50.33; H, 2.75; N, 5.10; Cl, 6.46; S, 5.84. Found: C, 50.42; H, 2.75; N, 5.01; Cl, 6.53; S, 5.76.


16.25



16.25

¹H-NMR (DMSO): δ 8.74 (s, 1H), 8.21 (m, 1H), 8.11 (d, J=1.48 Hz, 1H), 7.85 (m, 2H), 7.57 (d, J=8.44 Hz, 2H), 7.26 (m, 7H), 4.91 (s, 2H). MS (ES⁻): 513 (M-H, 100). Anal. Calc. for C₂₃H₁₆F₆N₂O₃S: C, 53.70; H, 3.13; N, 5.45; S, 6.23. Found: C, 53.65; H, 3.04; N, 5.33; S, 6.09.


16.26



16.26

¹H-NMR (DMSO): δ 8.75 (s, 1H), 8.21 (m, 1H), 8.08 (s, 1H), 7.83 (m, 2H), 7.57 (d, J=8.44 Hz, 2H), 7.27 (m, 2H), 7.15 (m, 2H), 6.81 (m, 2H), 4.82 (s, 2H), 3.67 (s, 3H). MS (ES⁻): 543 (M-H, 54). Anal. Calc. for C₂₄H₁₈F₆N₂O₄S: C, 52.94; H, 3.33; N, 5.15; S, 5.89. Found: C, 52.89; H, 3.37; N, 5.12; S, 5.97.


16.27



16.27

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.23 (m, 1H), 8.12 (d, J=1.76 Hz, 1H), 7.86 (m, 2H), 7.58 (d, J=8.40 Hz, 2H), 7.28 (m, 4H), 7.10 (m, 2H), 4.89 (s, 2H). MS (ES⁻): 531 (M-H, 100). Anal. Calc. for C₂₃H₁₅F₇N₂O₃S: C, 51.88; H, 2.84; N, 5.26; S, 6.02. Found: C, 51.91; H, 2.83; N, 5.25; S, 6.09.


16.28



16.28


¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.23 (m, 1H), 8.14 (d, J=1.72 Hz, 1H), 7.80 (m, 2H), 7.60 (d, J=8.32 Hz, 2H), 7.41 (m, 2H), 7.30 (m, 4H), 4.91 (s, 2H). MS (ES⁻): 593 (M-H, 100). Anal. Calc. for C₂₃H₁₅BrF₆N₂O₃S: C, 46.56; H, 2.55; N, 4.72; S, 5.40. Found: C, 46.71; H, 2.56; N, 4.67; S, 5.22.

5

16.29**16.29**

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.23 (m, 1H), 8.14 (d, J=1.28 Hz, 1H), 7.87 (m, 2H), 7.59 (d, J=8.40 Hz, 2H), 7.31 (m, 3H), 7.10 (m, 3H), 4.93 (s, 2H). MS (ES⁻): 531 (M-H, 100). Anal. Calc. for C₂₃H₁₅F₇N₂O₃S: C, 51.88; H, 2.84; N, 5.26; S, 6.02. Found: C, 51.89; H, 2.88; N, 5.23; S, 5.89.

10


16.30**16.30**

¹H-NMR (DMSO): δ 8.75 (s, 1H), 8.22 (m, 1H), 8.12 (s, 1H), 7.89 (d, J=8.08 Hz, 1H), 7.80 (m, 1H), 7.58 (d, J=8.40 Hz, 2H), 7.31 (m, 2H), 7.18 (m, 1H), 6.82 (d, J=7.68 Hz, 1H), 6.76 (m, 2H), 4.88 (s, 2H), 3.66 (s, 3H). MS (ES⁻): 543 (M-H, 54). Anal. Calc. for C₂₄H₁₈F₆N₂O₄S: C, 52.94; H, 3.33; N, 5.15; S, 5.89. Found: C, 53.15; H, 3.50; N, 5.15; S, 5.80.

15

20

16.31




16.31

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.23 (m, 1H), 8.09 (d, J=1.24 Hz, 1H), 7.90 (d, J=8.16 Hz, 1H), 7.80 (m, 1H), 7.57 (m, 3H), 7.46 (d, J=7.60 Hz, 1H), 7.30 (m, 3H), 7.17 (m, 1H), 4.97 (s, 2H). MS (ES⁻): 593 (M-H, 30). Anal. Calc. for C₂₃H₁₅BrF₆N₂O₃S: C, 46.56; H, 2.55; N, 4.72; S, 5.40. Found: C, 46.63; H, 2.57; N, 4.73; S, 5.36.

10

16.32




16.32

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.21 (m, 1H), 8.11 (s, 1H), 7.85 (m, 2H), 7.59 (m, 2H), 7.30 (m, 4H), 7.10 (m, 2H), 4.95 (s, 2H). MS (ES⁻): 531 (M-H, 100). Anal. Calc. for C₂₃H₁₅F₇N₂O₃S: C, 51.88; H, 2.84; N, 5.26; S, 6.02. Found: C, 51.83; H, 2.78; N, 5.16; S, 6.01.

15


16.33




16.33

20

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.23 (m, 1H), 8.12 (d, J=1.40 Hz, 1H), 7.80 (m, 2H), 7.58 (d, J=8.72 Hz, 2H), 7.47 (m, 2H), 7.30 (d, J=8.88 Hz, 2H), 7.22 (d, J=8.52 Hz, 2H), 4.89 (s, 2H). MS (ES⁻): 593 (M-H, 30). Anal. Calc. for C₂₃H₁₅BrF₆N₂O₃S: C, 46.56; H, 2.55; N, 4.72; S, 5.40. Found: C, 46.73; H, 2.51; N, 4.53; S, 5.31.


16.34**16.34**

¹H-NMR (DMSO): δ 8.82 (s, 1H), 8.15 (dd, J=1.64, 7.04 Hz, 1H), 7.95 (s, 1H), 7.76 (m, 2H), 7.66 (d, J=8.40 Hz, 2H), 7.27 (m, 5H), 7.11 (d, J=6.88 Hz, 2H), 3.92 (t, J=7.64 Hz, 2H), 2.67 (t, J=7.32 Hz, 2H). MS (ES⁻): 527 (M-H, 54). Anal. Calc. for C₂₄H₁₈F₆N₂O₃S: C, 54.55; H, 3.43; N, 5.30; S, 6.07. Found: C, 54.64; H, 3.37; N, 5.25; S, 6.10.

16.35**16.35**

¹H-NMR (DMSO): δ 8.76 (s, 1H), 8.20 (d, J=6.44 Hz, 1H), 8.08 (s, 1H), 7.82 (m, 2H), 7.59 (d, J=8.36 Hz, 2H), 7.29 (m, 2H), 6.71 (m, 3H), 4.78 (s, 2H), 4.17 (s, 4H). MS (ES⁻): 571 (M-H, 100). Anal. Calc. for C₂₅H₁₈F₆N₂O₅S: C, 52.45; H, 3.17; N, 4.89; S, 5.60. Found: C, 52.34; H, 3.13; N, 4.80; S, 5.46.


16.36



16.36

¹H-NMR (DMSO): δ 8.81 (s, 1H), 7.88 (s, 1H), 7.75 (m, 5H), 7.39 (m, 1H), 6.61 (m, 1H), 3.06 (s, 3H), 1.49 (s, 9H). MS (ES⁻): 468 (M-H, 100). Anal. Calc. for C₂₀H₂₁F₆NO₃S: C, 51.57; H, 4.51; N, 2.98; S, 6.83. Found: C, 51.14; H, 4.55; N, 2.96; S, 6.77.


16.37



16.37

¹H-NMR (DMSO): δ 9.75 (s, 1H), 8.19 (m, 1H), 8.098 (d, J=1.64 Hz, 1H), 7.91 (m, 1H), 7.80 (t, J=7.88 Hz, 1H), 7.58 (d, J=8.52 Hz, 2H), 7.25 (m, 4H), 6.86 (m, 2H), 4.84 (s, 2H), 3.55 (s, 3H). MS (ES⁻): 453 (M-H, 92). Anal. Calc. for C₂₄H₁₈F₆N₂O₄S: C, 52.94; H, 3.33; N, 5.15; S, 5.89. Found: C, 53.00; H, 3.21; N, 5.13; S, 5.91.

16.38



16.38


¹H NMR (CDCl₃): δ 7.74 (d, J= 8.7 Hz, 2H), 7.22 (d, J= 8.7 Hz, 2H), 6.93 (s, 1H), 4.12 (h, J = 6.7 Hz, 1H), 3.55 (s, 1H), 1.15 (d, J = 6.7 Hz, 6H). MS (EI)

m/z 515 (M^+). Anal. Calcd. for $C_{16}H_{13}Cl_2F_6NO_3S_2$: C, 37.22; H, 2.54; N, 2.71; S, 12.42. Found: C, 37.33; H, 2.47; N, 2.65; S, 12.48.

5

EXAMPLE 17

This example illustrates the preparation of compound 17.1


**17.1**

10 Nitric acid (90%, 0.2 mL) was added to compound 14.11 (273 mg, 0.5 mmol) in acetic acid (2 mL) at 0°C. The mixture was stirred at room temperature for 18 hour, at 60°C for 6 hours, and at 100°C for 3 hours. The mixture was poured into ice and extracted with EtOAc. The organic layer was washed with sodium bicarbonate solution and 0.1 N HCl, dried, and concentrated. The residue was
15 purified by silica column chromatography (eluted with 1:5 EtOAc/Hexane) to give 210 mg of product.

1H NMR ($CDCl_3$): δ 10.35 (s, 1H), 8.63 (s, 3H), 7.93 (d, $J=9$ Hz, 1H), 7.86 (d, $J=9$ Hz, 1H), 7.23 (s, 1H), 3.90 (s, 1H). MS (ES) m/z 516.9 ($[M-H]^+$). Anal. Calcd. for $C_{13}H_6Cl_2F_6N_2O_5S_2 \cdot 1/3EtOAc$: C, 31.38; H, 1.59; N, 5.11; S, 11.69. Found:
20 C, 31.53; H, 1.58; N, 5.09; S, 11.85.

EXAMPLE 18

This example illustrates the preparation of compound 18.1.

**18.1**

25

A mixture of N-methylaniline (1.08 mL, 10 mmol), pipsyl chloride (3.02 g, 10 mmol), and 2,6-lutidine (2.33 mL, 20 mmol) in acetone (30 mL) was stirred at room temperature for 20 hours. The lutidine•HCl salt was filtered. The filtrate was washed with hexane and water and dried to give 3.6 g of a white solid (N-methyl-N-phenyl-4-iodobenzenesulfonamide).

¹H NMR (CDCl₃): δ 7.80 (d, J= 8.5 Hz, 2H), 7.23-7.35 (m, 5H), 7.09 (d, J= 8.5 Hz, 2H), 3.18 (s, 3H). MS (EI) *m/z* 373 (M⁺).


n-BuLi (1.6 M, 1.25 mL, 2 mmol) was added to the above product (373 mg, 1 mmol) in THF (10 mL) at -78°C. The mixture was stirred for 30 minutes, and this mixture was added dropwise to a solution of methyl trifluoroacetate (0.4 mL, 4 mmol) in THF (10 mL) at -78°C. The mixture was allowed to warm to room temperature slowly and stirred at room temperature for 1 hour. The mixture was treated with EtOAc, and it was washed with brine, dried, and concentrated. The residue was purified by silica column chromatography (eluted with 30% EtOAc/Hexane) to give 200 mg of product (N-methyl-N-phenyl-4-trifluoroacetylbenzenesulfonamide).

¹H NMR (CDCl₃): δ 8.14 (d, J= 8.0 Hz, 2H), 7.72 (d, J= 8.7 Hz, 2H), 7.32 (m, 3H), 7.09 (d, J= 8.0 Hz, 2H), 3.24 (s, 3H). MS (ES) *m/z* 362.0 ([M+H₃O]⁺).


Tetrabutylammonium fluoride (1.0 M in THF, 0.01 mL) was added to a solution of the above product (186 mg, 0.54 mmol) and trifluoromethyl trimethylsilane (0.5 M in THF, 1.62 mL, 0.81 mmol) in THF (7 mL) at 0°C. Then the mixture was stirred at room temperature overnight. EtOAc (50 mL) was added to the reaction mixture, and it was washed with brine, dried, and concentrated. The residue was purified by silica column chromatography (eluted with 30% EtOAc/Hexane) to give 90 mg of product (18.1).

¹H NMR (CDCl₃): δ 7.83 (d, J= 8.4 Hz, 2H), 7.62 (d, J= 7.0 Hz, 2H), 7.30 (m, 3H), 7.07 (d, J= 8.4 Hz, 2H), 4.11 (bs, 1H), 3.20 (s, 3H). MS (ES) *m/z* 412.1 ([M-H]⁻). Anal. Calcd. for C₁₆H₁₃F₆NO₃S: C, 46.49; H, 3.17; N, 3.39. Found: C, 46.71; H, 3.23; N, 3.36.


Similar procedures to the one described for the synthesis of compound 18.1 were followed to produce compounds 18.2 to 18.15.

18.2**18.2**

¹H-NMR (DMSO): δ 9.08 (s, 1H), 7.80 (m, 4H), 7.58 (m, 1H), 7.20 (m, 1H), 7.09 (m, 2H), 3.78 (t, J=5.96 Hz, 2H), 2.40 (m, 2H), 1.61 (m, 2H). MS (ES⁻): 438 (M-H, 42). Anal. Calc. for C₁₈H₁₅F₆NO₃S: C, 49.20; H, 3.43; N, 3.19; S, 7.30. Found: C, 49.42; H, 3.58; N, 5.15; S, 7.23.

18.3**18.3**


¹H-NMR (DMSO): δ 9.10 (s, 1H), 7.90 (d, J=8.52 Hz, 2H), 7.71 (m, 2H), 7.43 (m, 2H), 7.14 (m, 2H), 3.17 (s, 3H). MS (ES⁻): 446 (M-H, 100). Anal. Calc. for C₁₆H₁₂ClF₆NO₃S: C, 42.92; H, 2.70; N, 3.13; Cl, 7.92; S, 7.16. Found: C, 43.04; H, 2.69; N, 3.17; Cl, 8.01; S, 7.20.

18.4**18.4**

¹H-NMR (DMSO): δ 9.09 (s, 1H), 7.90 (d, J=8.44 Hz, 2H), 7.70 (m, 2H), 6.96 (m, 2H), 6.88 (m, 2H), 3.74 (s, 3H), 3.13 (s, 3H). MS (ES⁻): 442 (M-H, 62).

Anal. Calc. for $C_{17}H_{15}F_6NO_4S$: C, 46.05; H, 3.41; N, 3.16; S, 7.23. Found: C, 46.15; H, 3.29; N, 3.17; S, 7.21.


18.5



18.5

1H -NMR (DMSO): δ 9.09 (s, 1H), 7.92 (s, 4H), 7.41 (m, 3H), 7.03 (m, 2H), 4.45 (m, 1H), 0.96 (d, $J=6.76$ Hz, 6H). MS (ES⁺): 442 (M+H, 100). Anal. Calc. for $C_{18}H_{17}F_6NO_3S$: C, 48.98; H, 3.88; N, 3.17; S, 7.26. Found: C, 49.07; H, 3.76; N, 3.16; S, 7.19.


18.6



18.6

1H -NMR (DMSO): δ 9.13 (s, 1H), 7.94 (d, $J=8.44$ Hz, 2H), 7.86 (d, $J=8.72$ Hz, 2H), 7.22 (m, 6H), 7.09 (m, 1H), 6.95 (m, 2H), 4.87 (s, 2H). MS (ES⁺): 508 (M+H, 75). Anal. Calc. for $C_{22}H_{16}F_7NO_3S$: C, 52.07; H, 3.18; N, 2.76; S, 6.32. Found: C, 52.14; H, 3.28; N, 2.77; S, 6.42.


18.7



18.7


¹H-NMR (DMSO): δ 9.13 (s, 1H), 7.93 (d, J=8.40 Hz, 2H), 7.84 (m, 2H), 7.60 (d, J=8.00 Hz, 1H), 7.52 (t, J=8.08 Hz, 1H), 7.43 (d, J=8.36 Hz, 1H), 7.24 (m, 6H), 4.93 (s, 2H).

MS (ES⁺): 558 (M+H, 35). Anal. Calc. for C₂₃H₁₆F₉NO₃S: C, 49.56; H, 2.89; N, 2.51; S, 5.75. Found: C, 49.74; H, 3.01; N, 2.55; S, 5.70.


18.8**18.8**

10 ¹H-NMR (DMSO): δ 9.11 (s, 1H), 7.90 (d, J=8.44 Hz, 2H), 7.71 (m, 3H), 7.62 (m, 1H), 7.50 (m, 1H), 7.25 (3, 1H), 3.24 (s, 3H). MS (ES⁺): 482 (M+H, 100). Anal. Calc. for C₁₇H₁₂F₉NO₃S: C, 42.42; H, 2.51; N, 2.91; S, 6.66. Found: C, 42.69; H, 2.59; N, 3.01; S, 6.38.

15

18.9**18.9**


20 ¹H-NMR (DMSO): δ 9.13 (s, 1H), 7.96 (m, 2H), 7.86 (m, 2H), 7.2 (m, 7H), 6.83 (m, 1H), 6.74 (m, 1H), 4.86 (s, 2H), 2.26 (s, 3H). MS (ES⁺): 536 (M+H, 100). Anal. Calc. for C₂₃H₁₉F₆NO₃S₂: C, 51.58; H, 3.58; N, 2.62; S, 11.98. Found: C, 52.30; H, 3.66; N, 2.71; S, 12.14.

18.10

18.10

¹H-NMR (DMSO): δ 9.11 (s, 1H), 7.91 (d, J=8.32 Hz, 2H), 7.74 (m, 2H), 7.28 (m, 1H), 7.18 (d, J=7.12 Hz, 1H), 6.80 (d, J=7.76 Hz, 1H), 6.80 (d, J=1.48 Hz, 1H), 3.18 (s, 3H), 2.33 (s, 3H). MS (ES⁺): 460 (M+H, 87). Anal. Calc. for C₁₇H₁₅F₆NO₃S₂: C, 44.44; H, 3.29; N, 3.05; S, 13.96. Found: C, 45.24; H, 3.36; N, 3.10; S, 14.26.


18.11



18.11

¹H-NMR (DMSO): δ 9.12 (s, 1H), 7.90 (d, J=8.44 Hz, 2H), 7.77 (d, J=7.28 Hz, 1H), 7.72 (d, J=8.56 Hz, 2H), 7.56 (m, 3H), 3.20 (s, 3H). MS (ES⁺): 439 (M+H, 60). Anal. Calc. for C₁₇H₁₂F₆N₂O₃S: C, 46.58; H, 2.76; N, 6.39; S, 7.32. Found: C, 46.58; H, 2.78; N, 6.32; S, 17.22.


18.12



18.12

¹H-NMR (DMSO): δ 9.10 (s, 1H), 7.92 (s, 4H), 7.57 (m, 1H), 7.49 (m, 1H), 7.19 (m, 1H), 6.81 (s, 1H), 4.51 (m, 1H), 0.99 (d, J=6.80 Hz, 6H). MS (ES⁺): 526 (M+H, 80). Anal. Calc. for C₁₉H₁₆F₉NO₄S: C, 43.44; H, 3.07; N, 2.67; S, 6.10. Found: C, 44.52; H, 3.25; N, 2.75; S, 2.03.


18.13



18.13

¹H-NMR (DMSO): δ 9.10 (s, 1H), 7.92 (s, 4H), 7.31 (m, 2H), 6.83 (d, J=7.60 Hz, 1H). 6.68 (s, 1H), 4.49 (m, 1H), 2.34 (s, 3H), 0.98 (d, J=6.68 Hz, 6H). MS (ES⁺): 488 (M+H, 100). Anal. Calc. for C₁₉H₁₉F₆NO₃S₂: C, 46.81; H, 3.93; N, 2.87; S, 13.16. Found: C, 47.17; H, 3.98; N, 2.89; S, 12.93.


18.14



18.14

¹H-NMR (DMSO): δ 9.07 (s, 1H), 7.88 (s, 4H), 7.41 (m, 4H), 7.17 (m, 2H), 7.00 (m, 2H), 6.83 (m, 1H), 6.51 (m, 1H), 4.43 (m, 1H), 0.93 (d, J=6.72 Hz, 6H). MS (ES⁺): 534 (M+H, 75). Anal. Calc. for C₂₄H₂₁F₆NO₄S: C, 54.03; H, 3.97; N, 2.63; S, 6.01. Found: C, 54.10; H, 3.93; N, 2.65; S, 6.05.


18.15



18.15

¹H-NMR (DMSO): δ 9.10 (s, 1H), 7.95 (s, 4H), 7.20 (m, 1H), 7.50 (m, 6H), 7.08 (m, 2H), 4.55 (m, 1H), 1.02 (d, J=6.68 Hz, 6H). MS (ES⁺): 518 (M+H, 100). Anal. Calc. for C₂₄H₂₁F₆NO₃S: C, 55.70; H, 4.09; N, 2.71; S, 6.20. Found: C, 56.26; H, 4.27; N, 2.70; S, 6.02.

Additional examples prepared by procedures similar to those described above:


18.16

5

18.16


$^1\text{H-NMR}$ (CDCl_3): δ 7.78 (d, $J = 8.3$ Hz, 2H), 7.27 (d, $J = 8.3$ Hz, 2H), 3.90 (bs, 1H), 3.26 (s, 3H), 2.20 (m, 1H), 1.40-1.70 (m, 8H), 1.15-1.35 (m, 2H). MS (ES-): 382 (M-H). Anal. Calcd. For $\text{C}_{17}\text{H}_{19}\text{F}_6\text{NO}_2$: C, 53.27; H, 5.00; N, 3.65. Found: C, 53.72; H, 5.21; N, 3.47.

10

18.17**18.17**


$^1\text{H-NMR}$ (CDCl_3): δ 7.76 (d, $J = 8.4$ Hz, 2H), 7.32 (s, $J = 8.4$ Hz, 2H), 3.60 (s, 1H), 3.22 (s, 3H), 1.84 (s, 3H), 1.73 (s, 6H), 1.57 (d, $J = 12.4$ Hz, 3H), 1.47 (d, $J = 12.4$ Hz, 3H). MS (ES-): 434 (M-H). Anal. Calcd. For $\text{C}_{21}\text{H}_{23}\text{F}_6\text{NO}_2$: C, 57.93; H, 5.32; N, 3.22. Found: C, 58.07; H, 5.33; N, 3.18.

15


18.18**18.18**

20

¹H-NMR (CD₃OD): δ 7.61 (d, J = 8.4 Hz, 2H), 7.25 (m, 7H), 3.49 (s, 3H). MS (ES⁺): 378 (M+ H), MS (ES⁻): 376 (M-H). Anal. Calcd. for C₁₇H₁₃F₆NO₂: C, 54.12; H, 3.47; N, 3.71. Found: C, 54.33; H, 3.41; N, 3.70.


18.19**18.19**

¹H-NMR (CD₃OD): δ 7.65 (d, J = 8.4 Hz, 2H), 7.26 (d, J = 8.7 Hz, 4H), 7.19 (s, 2H), 3.48 (s, 3H). MS (ES⁺): 412 (M+ H), MS (ES⁻): 410 (M-H). Anal. Calcd. for C₁₇H₁₂ClF₆NO₂: C, 49.59; H, 2.94; N, 3.40. Found: C, 49.72; H, 2.96; N, 3.40.

18.20**18.20**

¹H-NMR (CDCl₃): δ 8.12 (d, J = 9.1 Hz, 1H), 7.90 (s, 1H), 7.74 (d, J = 7.0 Hz, 1H), 7.69 (d, J = 6.8 Hz, 2H), 7.62 (dd, J = 9.1, 7.0 Hz, 1H), 7.20 (d, J = 6.8 Hz, 2H), 4.82 (s, 2H), 3.80 (s, 1H), 3.22 (s, 3H). MS (ES⁻): 491 (M-H). Anal. Calcd. For C₁₆H₁₄F₆N₂O₅S₂: C, 39.02; H, 2.87; N, 5.69; S, 13.02. Found: C, 39.52; H, 2.87; N, 5.44; S, 12.71.

18.21




18.21

¹H-NMR (CDCl₃): δ 7.74 (d, J = 7.9 Hz, 2H), 7.69 (d, J = 8.4 Hz, 2H),
 5 7.57 (t, J = 7.6 Hz, 1H), 7.47 (t, J = 7.6 Hz, 2H), 7.15 (d, J = 8.5 Hz, 2H), 4.61 (m,
 1H), 3.46 (s, 1H), 1.05(d, J = 6.8 Hz, 6H). MS (ES⁺): 442 (M⁺ H), MS (ES⁻): 440
 (M-H). Anal. Calcd. for C₁₈H₁₇F₆NO₃S : C, 48.98; H, 3.88; N, 3.17; S, 7.26. Found:
 C, 49.08; H, 3.82; N, 3.09; S, 7.27.

10


18.22



18.22

¹H-NMR (CD₃OD): δ 7.88 (d, J = 8.9 Hz, 1H), 7.68 (d, J = 8.2 Hz,
 15 2H), 7.65 (t, J = 7.6 Hz, 1H), 7.52 (t, 7.8 Hz, 2H), 7.25 (d, J = 8.6 Hz, 1H), 7.14 (s,
 1H), 3.92 (t, J = 4.6 Hz, 2H), 3.75 (t, J = 4.6 Hz, 2H). MS (ES⁻): 440 (M-H). Anal.
 Calcd. for C₁₇H₁₃F₆NO₄S : C, 46.26; H, 2.97; N, 3.17; S, 7.27. Found: C, 46.39; H,
 2.97; N, 3.18; S, 7.29.

18.23




18.23

¹H-NMR (CDCl₃): δ 7.94 (d, J = 9.0 Hz, 1H), 7.59 (d, J = 5.0 Hz, 1H),
 5 7.48 (d, J = 5.1 Hz, 1H), 7.27 (d, 9.5 Hz, 1H), 7.23 (s, 1H), 7.07 (t, J = 5.0 Hz, 1H),
 3.93 (t, J = 5.0 Hz, 2H), 3.86 (t, J = 4.9 Hz, 2H), 3.37 (s, 1H). MS (ES⁺): 448 (M⁺
 H), MS (ES⁻): 446 (M-H). Anal. Calcd. for C₁₅H₁₁F₆NO₄S₂: C, 40.27; H, 2.48; N,
 3.13; S, 14.34. Found: C, 40.10; H, 2.51; N, 3.12; S, 14.32.

10


18.24



18.24

¹H-NMR (CDCl₃): δ 7.98 (s, 1H), 7.85 (m, 3H), 7.61 (t, J = 7.9 Hz,
 15 1H), 7.29 (d, J = 8.8 Hz, 1H), 7.23 (s, 1H), 4.05 (s, 1H), 3.94 (t, J = 3.6 Hz, 2H), 3.90
 (t, J = 3.6 Hz, 2H). MS (ES⁻): 465 (M-H). Anal. Calcd. for C₁₈H₁₂F₆N₂O₄S: C, 46.36;
 H, 2.59; N, 6.01; S, 6.88. Found: C, 46.76; H, 2.72; N, 5.76; S, 6.52.

18.25




18.25

- 5 $^1\text{H-NMR}$ (CDCl_3): δ 7.71 (d, $J = 8.8$ Hz, 1H), 7.27 (s, 1H), 7.24 (d, $J = 8.8$ Hz, 1H), 6.99 (s, 1H), 4.12 (t, $J = 4.6$ Hz, 2H), 3.99 (t, $J = 4.2$ Hz, 2H) 3.33 (s, 1H). MS (ES $^-$): 514 (M-H). Anal. Calcd. for $\text{C}_{15}\text{H}_9\text{Cl}_2\text{F}_6\text{NO}_4\text{S}_2$: C, 34.90; H, 1.76; N, 2.71; S, 12.42. Found: C, 35.00; H, 1.66; N, 2.71; S, 12.55.

10


18.26



18.26

- 15 $^1\text{H-NMR}$ (CDCl_3): δ 7.73 (d, $J = 7.9$ Hz, 2H), 7.57 (t, $J = 7.9$ Hz, 1H), 7.48 (m, 4H), 7.39 (d, $J = 8.8$ Hz, 1H), 3.72 (bs, 2H), 3.44 (s, 1H), 2.42 (t, $J = 4.5$ Hz, 2H), 1.80 (m, 2H), 1.57 (m, 2H). MS (ES $^+$): 454 (M+H), MS (ES $^-$): 452 (M-H). Anal. Calcd. for $\text{C}_{19}\text{H}_{17}\text{F}_6\text{NO}_3\text{S}$: C, 50.33; H, 3.78; N, 3.09; S, 7.07. Found: C, 50.06; H, 3.69; N, 3.06; S, 7.22.

18.27




18.27

5 $^1\text{H-NMR}$ (CDCl_3): δ 8.02 (s, 1H), 7.91 (d, $J = 8.0$ Hz, 1H), 7.84 (d, $J = 7.7$ Hz, 1H), 7.61 (t, $J = 7.9$ Hz, 1H), 7.539 (d, $J = 8.5$ Hz, 1H), 7.51 (s, 1H) 7.31 (d, $J = 8.2$ Hz, 1H), 4.06 (s, 1H), 3.74 (bs, 2H), 2.42 (t, $J = 5.0$ Hz, 2H), 1.81 (q, $J = 5.7$ Hz, 2H), 1.60 (bs, 2H). MS (ES $^-$): 477 (M-H). Anal. Calcd. for $\text{C}_{20}\text{H}_{16}\text{F}_6\text{N}_2\text{O}_3\text{S}$: C, 50.21; H, 3.37; N, 5.86; S, 6.70. Found: C, 51.28; H, 3.75; N, 5.50; S, 6.39.

10

18.28



18.28


15 Reductive amination:

A mixture of 4-(hexafluoro-2-hydroxyisopropyl)aniline (1.0 g, 0.386 mol) and tetrahydro-4H-pyran-4-one (0.433 g, 0.433 mol, available from Aldrich Chemical Company) in methanol was stirred overnight. To this solution was added acetic acid (0.25 mL) followed by sodium cyanoborohydride (0.365 g, 0.006 mol).
 20 Stirring was continued for an additional 3 hr, at which time TLC analysis revealed completion of reaction. The reaction mixture was concentrated under vacuum. The residue was dissolved in ethyl acetate, washed with brine, dried, and concentrated. Purification of the crude product by flash chromatography on silica gel eluted with 4:1 to 2:1 hexane/EtOAc produced pure product (18.28) 1.06 g, 80.0%.

¹H-NMR (CDCl₃): δ 7.48 (d, J = 8.0 Hz, 2H), 6.61 (d, J = 8.0 Hz, 2H), 4.00 (m, 1H), 3.83 (s, 1H), 3.50 (t, J = 6.5 Hz, 4H), 3.40 (s, 1H), 1.52 (m, 4H). MS (ES⁺): 344 (M+H).

5

18.29




18.29

Starting from compound 18.28 and following the standard procedure for sulfonylation, compound 18.29 was obtained.

¹H-NMR (CDCl₃): δ 7.72 (d, J = 7.7 Hz, 2H), 7.66 (d, J = 8.1 Hz, 2H), 7.57 (t, J = 7.3 Hz, 1H), 7.47 (t, J = 7.7 Hz, 2H), 7.09 (d, J = 8.4 Hz, 1H), 3.72 (m, 1H), 4.19 (s, 1H), 3.91 (dd, J = 11.5, 4 Hz, 2H), 3.45 (t, J = 11.8 Hz, 2H), 1.72 (d, J = 12 Hz, 2H), 1.44 (q, J = 12 Hz, 2H). MS (ES⁺): 484 (M+ H), MS (ES⁻): 482 (M-H).
 Anal. Calcd. for C₂₀H₁₉F₆NO₄S : C, 49.69; H, 3.96; N, 2.90; S, 6.63. Found: C, 49.66; H, 3.84; N, 2.89; S, 6.52.

15

18.30




18.30

Substituting cyclopentanone for tetrahydro-4H-pyran-4-one, compound 18.30 was prepared by a procedure similar to the synthesis of compound 18.28.

20

¹H-NMR (CDCl₃): δ 7.45 (d, J = 8.0 Hz, 2H), 6.60 (d, J = 8.0 Hz, 2H), 3.88 (s, 1H), 3.80 (s, 1H), 2.05 (m, 4H), 1.65 (m, 4H). MS (ES⁺): 260 (M+ H).

18.31




18.31

5 ¹H-NMR (CDCl₃): δ 7.70 (d, J=8.5 Hz, 2H), 7.59 (d, J=4.9 Hz, 1H), 7.43 (d, J=5.0 Hz, 1H), 7.19(d, 8.4 Hz, 2H), 7.06 (t, J=4.0 Hz, 1H), 4.52 (m, 1H), 3.45 (s, 1H), 1.89(t, J=5.2 Hz, 2H), 1.48 (m, 4H), 1.32 (m, 2H). MS (ES⁻): 472 (M-H).
 Anal. Calcd. for C₁₈H₁₇F₆NO₃S₂ : C, 45.66; H, 3.62; N, 2.96; S, 13.55. Found: C, 45.77; H, 3.69; N, 2.97; S, 13.69.

10


18.32



18.32


15 ¹H-NMR (CDCl₃): δ 8.01 (s, 1H), 7.89 (d, J=8.0 Hz, 2H), 7.85 (d, J=7.8 Hz, 1H), 7.71 (d, J=8.4 Hz, 2H), 7.61(t, J=7.9 Hz, 1H), 7.10 (d, J=8.7 Hz, 2H), 4.53 (m, 1H), 3.53 (s, 1H), 1.86(m, 2H), 1.51 (m, 4H), 1.30 (m, 2H). MS (ES⁻): 491 (M-H). Anal. Calcd. for C₂₁H₁₈F₆N₂O₃S : C, 51.22; H, 3.68; N, 5.69; S, 6.51. Found: C, 51.29; H, 3.56; N, 5.63; S, 6.49.

18.33



18.33

18.34



18.34


5

Compound **18.33** was prepared from the corresponding sulfonyl chloride following the standard sulfonylation procedure described above. Compound **18.33** (150 mg, 0.342 mmol) was treated under Lemieux-Johnson oxidative cleavage conditions (OsO_4 , 0.0342 mmol; NaIO_4 , 250 mg, 1.27 mmol; in 2:1 acetone/ H_2O) to give the corresponding aldehyde. The aldehyde was then reduced with NaBH_4 in methanol at 0°C to give compound **18.34**, 50 mg, 30%.

$^1\text{H-NMR}$ (CDCl_3): δ 7.82 (d, $J = 7.2$ Hz, 1H), 7.80 (bs, 1H), 7.55 (d, $J = 8.4$ Hz, 2H), 7.51 (t, $J = 7.2$ Hz, 1H), 7.37 (d, $J = 7.6$ Hz, 1H), 7.25 (t, $J = 7.6$ Hz, 1H), 7.13 (d, $J = 8.4$ Hz, 2H), 4.04 (t, $J = 6.1$ Hz, 2H), 3.57 (s, 1H), 3.43 (t, $J = 6.1$ Hz, 2H), 2.15 (bs, 1H). MS (ES $^-$): 442 (M $-$ H).

15

18.35



18.35

20

A sample of compound **18.34** (22 mg, 0.049 mmol) was dissolved in anhydrous THF (2 mL). To it was added triphenylphosphine (32.5 mg, 0.124 mmol)

followed by diethyl azodicarboxylate (15.6 mL, 0.10 mmol) at 0°C. Stirring was continued for an additional 2 hrs. The reaction mixture was diluted with ethyl acetate, washed with aqueous NaHCO₃ solution, dried over MgSO₄, filtered and concentrated.


Crude product was purified by flash chromatography on silica gel, eluted with 1:1

5 hexane/EtOAc to give compound **18.35**, 18.2 mg, 86.2%.

¹H-NMR (CDCl₃): δ 7.89 (d, J = 7.6 Hz, 1H), 7.69 (d, J = 8.4 Hz, 2H), 7.532 (t, J = 7.6 Hz, 1H), 7.45 (d, J = 7.6 Hz, 1H), 7.37 (d, J = 8.4 Hz, 2H), 7.26 (d, J = 7.6 Hz, 1H), 7.37 (d, J = 8.4 Hz, 2H), 7.26 (d, J = 7.6 Hz, 1H), 4.27 (t, J = 6.4 Hz, 2H), 3.58 (s, 1H), 3.19 (t, J = 6.4 Hz, 2H). MS (ES⁻): 424 (M⁻ H).

10

18.36



18.36

15 4-(Hexafluoro-2-hydroxyisopropyl)benzoic acid (100 mg, 0.35 mmol, available from Lancaster) was treated with oxalyl chloride (2.0M solution in dichloromethane, 0.35 mL, 0.70 mmol) in dichloromethane (2 mL) at ambient temperature for two hours. The reaction mixture was evaporated to dryness, redissolved in dichloromethane (2 mL), and was added to a stirred mixture of N-

20 methylaniline (56 mg, 0.53 mmol) and diisopropylethylamine (0.12 mL, 0.70 mmol) in dichloromethane (2 mL). Upon completion of reaction, the mixture was diluted with dichloromethane, washed with 0.5 N HCl, brine, and NaHCO₃ solution. The organic phase was dried over MgSO₄, filtered and concentrated. The crude product was purified by flash chromatography on silica gel to give compound **18.36**, 118 mg,

25 90.1%.

¹H-NMR (CDCl₃): δ 7.50 (d, J = 8.0 Hz, 2H), 7.30 (d, J = 7.0 Hz, 2H), 7.20 (t, J = 7.0 Hz, 2H), 7.14 (t, J = 7.0 Hz, 1H), 7.01 (d, J = 8.0 Hz, 2H), 3.80 (s, 1H), 3.50 (s, 3H). MS (ES⁻): 376 (M⁻ H).

EXAMPLE 19

LXR α agonist activity can be characterized in a cell-based assay or a
5 peptide-sensor assay. The same assays can be utilized to characterize LXR α
antagonist activity, by evaluating the ability of a compound to block the effects of an LXR α
agonist, such as 24,25-epoxycholesterol.

Briefly, for the cell-based assay, a DNA-binding domain of the
nonreceptor transcription factor GAL4 was fused to the putative ligand-binding
10 domain of LXR α . The resultant construct was introduced into 293 cells, together with
UAS-containing luciferase reporter construct. The transfected cells were then treated
with chemical libraries and luciferase activity was measured. Individual compounds
were evaluated relative to a control (no addition of compound) and the EC₅₀ was
determined as the concentration necessary to produce 50% of the maximal luciferase
15 activity.


The peptide sensor LXR assay is an *in vitro* assay and was conducted
as generally described in co-pending applications Ser. Nos. 08/975,614 (filed
November 21, 1997) and 09/163,713 (filed September 30, 1998). Exemplary of the
activity observed for the compounds provided herein was exhibited by compound 2
20 (EC₅₀ in each of the assays was less than 2 micromolar).

All publications and patent applications cited in this specification are
herein incorporated by reference as if each individual publication or patent application
were specifically and individually indicated to be incorporated by reference.
25 Although the foregoing invention has been described in some detail by way of
illustration and example for purposes of clarity of understanding, it will be readily
apparent to those of ordinary skill in the art in light of the teachings of this invention
that certain changes and modifications may be made thereto without departing from
the spirit or scope of the appended claims.

30

WHAT IS CLAIMED IS:

1. A composition for modulation of LXR function in a cell, said composition comprising a pharmaceutically acceptable excipient and a compound
 5 having the formula:



or a pharmaceutically acceptable salt thereof, wherein

Ar is an aryl group;

R¹ is a member selected from the group consisting of -OH, -CO₂H, -O-(C₁-C₇)alkyl, -OC(O)-(C₁-C₇)alkyl, -O-(C₁-C₇)heteroalkyl, -OC(O)-(C₁-C₇)heteroalkyl,
 10 -NH₂, -NH(C₁-C₇)alkyl, -N((C₁-C₇)alkyl)₂ and -NH-S(O)₂-(C₁-C₃)alkyl;

R² is a member selected from the group consisting of (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl;

X¹, X², X³, X⁴, X⁵ and X⁶ are each independently a member selected from the group consisting of H, (C₁-C₃)alkyl, (C₁-C₃)heteroalkyl, F and Cl, with the
 15 proviso that no more than three of X¹ through X⁶ are H, (C₁-C₃)alkyl or (C₁-C₇)heteroalkyl; and

Y is a divalent linking group selected from the group consisting of -N(R¹²)S(O)_m-, -N(R¹²)S(O)_mN(R¹³)-, -N(R¹²)C(O)-, -N(R¹²)C(O)N(R¹³)-,
 20 -N(R¹²)C(S)- and -N(R¹²)C(O)O-;

wherein R¹² and R¹³ are each independently selected from the group consisting of H, (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl, and optionally when Y is -N(R¹²)S(O)_m- or -N(R¹²)S(O)_mN(R¹³)-, R¹² forms a five- or six-membered ring fused to Ar or to R² through covalent attachment to Ar or to R²,
 respectively; and the subscript m is an integer of from 1 to 2;

with the proviso that when R¹ is OH, and -Y-R² is -N(R¹²)S(O)_m-R² or -N(R¹²)C(O)N(R¹³)-R² and is attached to a position para to the quaternary carbon attached to Ar, and when R² is phenyl, benzyl or benzoyl, then i) at least one of R¹² or R¹³ is other than hydrogen and contains an electron-withdrawing substituent, or ii) R² is substituted with a moiety other than amino, acetamido, di(C₁-C₇)alkylamino, (C₁-
 30 C₇)alkylamino, halogen, hydroxy, nitro, or (C₁-C₇)alkyl, or iii) the benzene ring

portion of R^2 is substituted with at least three independently selected groups in addition to the Y group or point of attachment to Y.

2. A composition in accordance with claim 1, wherein Ar represents a substituted or unsubstituted ring selected from the group consisting of benzene, naphthalene, pyridine, quinoline, isoquinoline, pyrrole, furan and thiophene.

3. A composition in accordance with claim 1, wherein Ar represents a benzene ring.


4. A composition in accordance with claim 1, wherein R^1 is a member selected from the group consisting of -OH, -CO₂H, -NH₂, -NH(C₁-C₇)alkyl, -N((C₁-C₇)alkyl)₂ and -NH-S(O)₂CH₃.

5. A composition in accordance with claim 1, wherein R^2 is aryl.

6. A composition in accordance with claim 1, wherein R^2 is phenyl, said phenyl group being substituted with at least one substituent at the ortho or meta position relative to the point of attachment to Y.


7. A composition in accordance with claim 1, wherein no more than one of X^1 through X^6 is hydrogen or (C₁-C₅)alkyl.

8. A composition in accordance with claim 1, selected from the group consisting of




wherein the subscript n represents an integer of from 0 to 4; and each R^{11} is independently selected from the group consisting of -OH, -NH₂, lower alkyl, lower alkoxy, -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR''C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -CN and -NO₂

9. A composition in accordance with claim 1, said compound having a formula selected from the group consisting of




10. A composition in accordance with claim 9, said compound having a formula selected from the group consisting of



5 11. A composition in accordance with claim 1, wherein X^1 , X^2 , X^3 , X^4 , X^5 and X^6 are each F.


12. A composition in accordance with claim 1, wherein X^1 , X^2 , X^3 , X^4 , X^5 and X^6 are each F, R^1 is a member selected from the group consisting of -OH, -NH₂, -CO₂H, -O-(C₁-C₇)alkyl, and -OC(O)-(C₁-C₇)alkyl, Y is a member selected from the group consisting of -N(R¹²)S(O)_m-, -N(R¹²)C(O)- and -N(R¹²)C(O)N(R¹³)-, and R² is aryl.

13. A composition in accordance with claim 10, said compound having a formula selected from the group consisting of:




wherein R^1 is a member selected from the group consisting of $-OH$ and $-NH_2$; X^1 and X^6 are each independently hydrogen or fluorine, R^{12} is fluoro(C_1 - C_4)alkyl; and R^2 is aryl.

14. A composition in accordance with claim 10, said compound
5 having a formula selected from the group consisting of:



wherein R^1 is a member selected from the group consisting of $-OH$ and $-NH_2$; X^1 and X^6 are each independently hydrogen or fluorine, R^{12} is H , (C_1 - C_4)alkyl or (C_2 - C_4)heteroalkyl; and R^2 is substituted or unsubstituted thienyl.


15. A composition in accordance with claim 10, said compound
10 having a formula selected from the group consisting of:




- wherein R^1 is a member selected from the group consisting of $-OH$ and $-NH_2$; X^1 and X^6 are each independently H or F ; R^{12} is (C_1 - C_4)alkyl; and R^2 is phenyl substituted with at least one member selected from the group consisting of halogen $-CN$, $-CF_3$, $-O-(C_1-C_4)alkyl$, $-C(O)-(C_1-C_4)alkyl$, $-C(O)-O(C_1-C_4)alkyl$, $-C(O)-NH(C_1-C_4)alkyl$ and $-C(O)N((C_1-C_4)alkyl)_2$.
15

16. A composition in accordance with claim 15, wherein R^1 is $-OH$; X^1 and X^6 are each fluorine; and R^2 is phenyl substituted with at least one member selected from the group consisting of $-CN$, $-CF_3$ and $-O-(C_1-C_4)alkyl$.


17. A composition in accordance with claim 1, wherein said compound is selected from the group consisting of:




18. A composition in accordance with claim 1, wherein said compound is selected from the group consisting of:



5 19. A composition in accordance with claim 1, wherein said compound is selected from the group consisting of:



20. A compound having the formula:



or a pharmaceutically acceptable salt thereof, wherein

Ar is an aryl group;

R¹ is a member selected from the group consisting of -OH, -CO₂H, -O-(C₁-C₇)alkyl, -OC(O)-(C₁-C₇)alkyl, -O-(C₁-C₇)heteroalkyl, -OC(O)-(C₁-C₇)heteroalkyl, -NH₂, -NH(C₁-C₇)alkyl, -N((C₁-C₇)alkyl)₂ and -NH-S(O)₂-(C₁-C₃)alkyl ;

R² is a member selected from the group consisting of (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl;

X¹, X², X³, X⁴, X⁵ and X⁶ are each independently a member selected from the group consisting of H, (C₁-C₃)alkyl, (C₁-C₃)heteroalkyl, F and Cl, with the proviso that no more than three of X¹ through X⁶ are H, (C₁-C₃)alkyl or (C₁-C₃)heteroalkyl; and

Y is a divalent linking group selected from the group consisting of -N(R¹²)S(O)_m-, -N(R¹²)S(O)_mN(R¹³)-, -N(R¹²)C(O)-, -N(R¹²)C(O)N(R¹³)-, -N(R¹²)C(S)- and -N(R¹²)C(O)O-;

wherein R¹² and R¹³ are each independently selected from the group consisting of H, (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl, and optionally when Y is -N(R¹²)S(O)_m- or -N(R¹²)S(O)_mN(R¹³)-, R¹² forms a five- or six-membered ring fused to Ar or to R² through covalent attachment to Ar or to R², respectively; and the subscript m is an integer of from 1 to 2;

with the proviso that when R¹ is OH, and -Y-R² is -N(R¹²)S(O)_m-R² or -N(R¹²)C(O)N(R¹³)-R² and is attached to a position para to the quaternary carbon attached to Ar, and when R² is phenyl, benzyl or benzoyl, then i) at least one of R¹² or R¹³ is other than hydrogen and contains an electron-withdrawing substituent, or ii) R² is substituted with a moiety other than amino, acetamido, di(C₁-C₇)alkylamino, (C₁-C₇)alkylamino, halogen, hydroxy, nitro, or (C₁-C₇)alkyl, or iii) the benzene ring portion of R² is trisubstituted in addition to the attached Y group.

21. A compound in accordance with claim 20, wherein Ar represents a substituted or unsubstituted ring selected from the group consisting of benzene, naphthalene, pyridine, quinoline, isoquinoline, pyrrole, furan and thiophene.

22. A compound in accordance with claim 20, wherein Ar represents a benzene ring.


23. A compound in accordance with claim 20, wherein R¹ is a member selected from the group consisting of -OH, -CO₂H, -NH₂, -NH(C₁-C₇)alkyl, -N((C₁-C₇)alkyl)₂ and -NH-S(O)₂CH₃.

24. A compound in accordance with claim 20, wherein R^2 is aryl.

25. A compound in accordance with claim 20, wherein R^2 is phenyl, said phenyl group being substituted with at least one substituent at the ortho or meta position relative to the point of attachment to Y.

5 26. A compound in accordance with claim 20, wherein no more than one of X^1 through X^6 is hydrogen or (C_1-C_5) alkyl.

27. A compound in accordance with claim 20, selected from the group consisting of




wherein the subscript n represents an integer of from 0 to 4; and


10 each R^{11} is independently selected from the group consisting of -OH, -NH₂, lower alkyl, lower alkoxy, -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)NR'R''', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -CN and -NO₂.

15 wherein R', R'' and R''' are each independently selected from the group consisting of (C_1-C_5) alkyl and (C_1-C_5) haloalkyl.

28. A compound in accordance with claim 27, said compound having a formula selected from the group consisting of:




29. A compound in accordance with claim 28, said compound having a formula selected from the group consisting of



30. A compound in accordance with claim 20, wherein X^1 , X^2 , X^3 , X^4 , X^5 and X^6 are each F.


31. A compound in accordance with claim 20, wherein X^1 , X^2 , X^3 , X^4 , X^5 and X^6 are each F; R^1 is a member selected from the group consisting of -OH, -CO₂H, -NH₂, -O-(C₁-C₇)alkyl, and -OC(O)-(C₁-C₇)alkyl; Y is a member selected from the group consisting of -N(R¹²)S(O)_m-, -N(R¹²)C(O)- and -N(R¹²)C(O)N(R¹³)-; and R² is aryl.

32. A compound in accordance with claim 29, said compound having the formula:




wherein R^1 is a member selected from the group consisting of $-OH$ and $-NH_2$; X^1 and X^6 are each independently hydrogen or fluorine, R^{12} is fluoro(C_1 - C_4)alkyl; and R^2 is aryl.

33. A compound in accordance with claim 29, said compound having
5 the formula:



wherein R^1 is a member selected from the group consisting of $-OH$ and $-NH_2$; X^1 and X^6 are each independently H or F; R^{12} is H, (C_1 - C_4)alkyl or (C_2 - C_4)heteroalkyl; and R^2 is substituted or unsubstituted thienyl.


34. A compound in accordance with claim 29, said compound having
10 the formula:




- wherein R^1 is a member selected from the group consisting of $-OH$ and $-NH_2$; X^1 and X^6 are each independently hydrogen or fluorine, R^{12} is (C_1 - C_4)alkyl; and R^2 is phenyl substituted with at least one member selected from the group consisting of halogen, $-CN$, $-CF_3$, $-O-(C_1-C_4)alkyl$, $-C(O)-(C_1-C_4)alkyl$, $-C(O)-O(C_1-C_4)alkyl$,
15 $-C(O)-NH(C_1-C_4)alkyl$ and $-C(O)N((C_1-C_4)alkyl)_2$.

35. A compound in accordance with claim 34, wherein R^1 is $-OH$; X^1 and X^6 are each fluorine; and R^2 is phenyl substituted with at least one member selected from the group consisting of $-CN$, $-CF_3$, and $-O-(C_1-C_4)alkyl$.


36. A compound in accordance with claim 20, wherein said compound is selected from the group consisting of:




37. A compound in accordance with claim 20, wherein said compound is selected from the group consisting of:



38. A compound in accordance with claim 20, wherein said compound is selected from the group consisting of:



40. A method of modulating LXR function in a cell, tissue or animal, said method comprising contacting said cell, tissue or animal with a pharmaceutical composition comprising an LXR-modulating amount of a compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein

Ar is an aryl group;

R¹ is a member selected from the group consisting of -OH, -CO₂H, -O-(C₁-C₇)alkyl, -OC(O)-(C₁-C₇)alkyl, -O-(C₁-C₇)heteroalkyl, -OC(O)-(C₁-C₇)heteroalkyl, -NH₂, -NH(C₁-C₇)alkyl, -N((C₁-C₇)alkyl)₂ and -NH-S(O)₂-(C₁-C₃)alkyl ;

R² is a member selected from the group consisting of (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl;

X¹, X², X³, X⁴, X⁵ and X⁶ are each independently a member selected from the group consisting of H, (C₁-C₃)alkyl, (C₁-C₃)heteroalkyl, F and Cl, with the proviso that no more than three of X¹ through X⁶ are H, (C₁-C₃)alkyl or (C₁-C₃)heteroalkyl; and

Y is a member selected from the group consisting of -N(R¹²)S(O)_m-, -N(R¹²)S(O)_mN(R¹³)-, -N(R¹²)C(O)-, -N(R¹²)C(O)N(R¹³)-, -N(R¹²)C(S)- and -N(R¹²)C(O)O-;

wherein R¹² and R¹³ are each independently selected from the group consisting of H, (C₁-C₇)alkyl, (C₁-C₇)heteroalkyl, aryl and aryl(C₁-C₇)alkyl, and optionally when Y is -N(R¹²)S(O)_m- or -N(R¹²)S(O)_mN(R¹³)-, R¹² forms a five- or six-membered ring fused to Ar or to R² through covalent attachment to Ar or to R², respectively; and the subscript m is an integer of from 1 to 2.

41. A method in accordance with claim 40, wherein said LXR function is associated with a disease or condition selected from the group consisting of lipid disorders and other metabolic disorders.


42. A method in accordance with claim 41, wherein said disease or condition is selected from the group consisting of atherosclerosis, elevated LDL plasma levels, low HDL plasma levels, hypertriglyceridemia, hypertension and hypercholesterolemia.

43. A method in accordance with claim 41, wherein said disease or condition is selected from the group consisting of obesity, diabetes, vitamin deficiency, osteoporosis, and multidrug resistance to chemotherapeutic drugs.


44. A method in accordance with claim 40, wherein said compound is administered in combination with a second lipid-lowering or cholesterol-lowering agent.

45. A composition of claim 1, further comprising a second lipid-lowering or cholesterol-lowering agent.


46. A composition in accordance with claim 1, wherein said compound has the formula:



5 47. A composition in accordance with claim 1, wherein said compound has the formula:



48. A composition in accordance with claim 1, wherein said compound has the formula:






FIG. 1.

2/2




FIG. 2.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 September 2000 (21.09.2000)

PCT

(10) International Publication Number
WO 00/54759 A3

- (51) International Patent Classification⁷: A61K 31/18, 31/277, 31/24, 31/381, 31/27, 31/17, 31/167, C07C 311/21, C07D 333/34
- (74) Agents: KEZER, William, B. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111 (US).
- (21) International Application Number: PCT/US00/06611
- (22) International Filing Date: 15 March 2000 (15.03.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/124,525 15 March 1999 (15.03.1999) US
- (71) Applicant: TULARIK INC. [US/US]; Two Corporate Drive, South San Francisco, CA 94080 (US).
- (72) Inventors: LI, Leping; 2100 Davis Drive, Burlingame, CA 94010 (US). MEDINA, Julio, C.; 1407 Cedar St., San Carlos, CA 94070 (US). HASEGAWA, Hirohiko; 1-98, 1 Kasugadenaka 3-chome, Konohana-ku, Osaka 554 (JP). CUTLER, Serena, T.; 449 Hawthorne Ave. #3, Palo Alto, CA 94301 (US). LIU, Jiwen; Apt. 211, 1075 Rollins Rd., Burlingame, CA 94010 (US). ZHU, Liusheng; Apt. 211, 2001 Trousdale Drive, Burlingame, CA 94010 (US). SHAN, Bei; 349 Quay Lane, Redwood City, CA 94065 (US). LUSTIG, Kevin; Two Corporate Drive, South San Francisco, CA 94080 (US).
- (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— With international search report.
- (88) Date of publication of the international search report:
15 February 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/54759 A3

(54) Title: LXR MODULATORS

(57) Abstract: The invention provides compounds, compositions and methods for modulating the effects of LXR in a cell. The compounds and compositions are useful both as diagnostic indicators of LXR function and as pharmacologically active agents. The compounds and compositions find particular use in the treatment of disease states associated with cholesterol metabolism, particularly atherosclerosis and hypercholesterolemia.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 00/06611

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/18 A61K31/277 A61K31/24 A61K31/381 A61K31/27
A61K31/17 A61K31/167 C07C311/21 C07D333/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols).

IPC 7 A61K C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, MEDLINE, BIOSIS, EMBASE, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	<p>WO 00 46203 A (MILLENNIUM PHARM INC) 10 August 2000 (2000-08-10)</p> <p>abstract page 9, line 1 -page 11, line 18 page 27 page 47 -page 48; claims --- -/--</p>	<p>1-5, 7-9, 11, 12, 20-24, 26-28, 30, 31, 40-45</p>



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

26 October 2000

Date of mailing of the international search report

08/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoff, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/06611

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	EP 0 919 542 A (NIHON NOHYAKU CO LTD) 2 June 1999 (1999-06-02) abstract page 71 -page 72 page 75 page 80; claims	1-9, 11, 12, 20-28, 30, 31
X	US 4 230 635 A (NEUSTADT BERNARD R) 28 October 1980 (1980-10-28) the whole document	1-13, 20-32, 40-42, 46-48
X	US 4 093 742 A (NEUSTADT BERNARD R) 6 June 1978 (1978-06-06) the whole document	1-4, 7-9, 11, 20-23, 26-28, 30, 40-42
X	GB 1 507 340 A (SCHERICO LTD) 12 April 1978 (1978-04-12) the whole document	1-4, 7-9, 11, 20-23, 26-28, 30, 40-42
X	US 3 281 466 A (H.C. STECKER) 25 October 1966 (1966-10-25) the whole document	1-9, 11, 12, 20-28, 30, 31
X	DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; MIRYAN, N. I. ET AL: "Derivatives of pyridinecarboxylic acids. Synthesis and antiexudative effect of fluorinated derivatives of nicotinamide and isonicotinamide" retrieved from STN Database accession no. 86:139798 CA XP002151153 abstract & KHIM.-FARM. ZH. (1977), 11(1), 70-2 , 1977,	1-5, 7-9, 11, 12, 20-24, 26-28, 30, 31

-/--

INTERNATIONAL SEARCH REPORT

Internr al Application No

PCT/US 00/06611

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; POLISHCHUK, V. R. ET AL: "Electron paramagnetic resonance spectra of 2-arylpolyfluoroisopropyl radicals" retrieved from STN Database accession no. 91:4737 CA XP002151154 abstract & IZV. AKAD. NAUK SSSR, SER. KHIM. (1979), (3), 659-61 , 1979,</p>	<p>20-23, 26-28,30</p>
X	<p>DATABASE 'Online! GILBERT, EVERETT E. ET AL: "perhalo ketones - (VI) aromatic amino derivs. of the penhaloacetones" retrieved from STN Database accession no. 16091f XP002151155 abstract</p>	<p>20-23, 26-28,30</p>
X	<p>US 4 251 534 A (ALDRICH PAUL E ET AL) 17 February 1981 (1981-02-17) examples 22-33</p>	<p>20-23, 26-28,30</p>
A	<p>US 4 240 979 A (ADOLPHI HEINRICH ET AL) 23 December 1980 (1980-12-23) the whole document</p>	<p>1-9,11, 12,20-28</p>
A	<p>EP 0 193 249 A (DUPHAR INT RES) 3 September 1986 (1986-09-03) abstract; table A compound no.4 claims</p>	<p>1-9,11, 12,20-28</p>

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 1-9,11,12,20-28,30,31,40-48 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds prepared in the examples and closely related homologous compounds, with due regard to the general idea underlying the present invention.

Claims searched completely: 10,13-19,29,32-38

Claims searched incompletely: 1-9,11,12,20-28,30,31,40-48

Remark: claim 39 is missing

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/06611

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0046203	A	10-08-2000	NONE	
EP 0919542	A	02-06-1999	AU 712421 B	04-11-1999
			AU 9329298 A	24-06-1999
			BR 9805060 A	21-03-2000
			CZ 9803799 A	16-06-1999
			HU 9802725 A	30-08-1999
			JP 11240857 A	07-09-1999
			PL 329898 A	07-06-1999
			ZA 9810677 A	26-05-1999
US 4230635	A	28-10-1980	NONE	
US 4093742	A	06-06-1978	ES 442992 A	01-08-1977
			ES 457806 A	01-03-1978
			ES 457807 A	01-03-1978
			US 4081547 A	28-03-1978
			US 4058612 A	15-11-1977
			US 4159335 A	26-06-1979
			AR 219478 A	29-08-1980
			AR 218436 A	13-06-1980
			AT 368146 B	10-09-1982
			AT 520678 A	15-01-1982
			AT 357557 B	25-07-1980
			AT 898575 A	15-12-1979
			AU 4416979 A	31-05-1979
			AU 501512 B	21-06-1979
			AU 8698975 A	02-06-1977
			BE 835899 A	25-05-1976
			CA 1063112 A	25-09-1979
			CH 623044 A	15-05-1981
			CS 188299 B	28-02-1979
			CS 188259 B	28-02-1979
			DD 124730 A	09-03-1977
			DE 2552933 A	10-06-1976
			DK 535275 A	03-06-1976
			FI 753332 A	03-06-1976
			FR 2346000 A	28-10-1977
			GB 1507340 A	12-04-1978
			HU 175541 B	28-08-1980
			IE 42759 B	08-10-1980
			IL 48557 A	31-05-1979
			IT 1051024 B	21-04-1981
			JP 51082239 A	19-07-1976
			LU 73888 A	06-09-1976
			NL 7513804 A	04-06-1976
			NO 753984 A,B,	03-06-1976
			PH 14474 A	07-08-1981
			SE 7513331 A	03-06-1976
			SE 8004364 A	11-06-1980
			SU 655311 A	30-03-1979
			SU 685146 A	05-09-1979
			ZA 7507457 A	24-11-1976
GB 1507340	A	12-04-1978	AR 219478 A	29-08-1980
			AR 218436 A	13-06-1980
			AT 368146 B	10-09-1982
			AT 520678 A	15-01-1982

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/06611

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1507340 A		AT 357557 B	25-07-1980
		AT 898575 A	15-12-1979
		AU 4416979 A	31-05-1979
		AU 501512 B	21-06-1979
		AU 8698975 A	02-06-1977
		BE 835899 A	25-05-1976
		CA 1063112 A	25-09-1979
		CH 623044 A	15-05-1981
		CS 188299 B	28-02-1979
		CS 188259 B	28-02-1979
		DD 124730 A	09-03-1977
		DE 2552933 A	10-06-1976
		DK 535275 A	03-06-1976
		ES 442992 A	01-08-1977
		ES 457806 A	01-03-1978
		ES 457807 A	01-03-1978
		FI 753332 A	03-06-1976
		FR 2346000 A	28-10-1977
		HU 175541 B	28-08-1980
		IE 42759 B	08-10-1980
		IL 48557 A	31-05-1979
		IT 1051024 B	21-04-1981
		JP 51082239 A	19-07-1976
		LU 73888 A	06-09-1976
		NL 7513804 A	04-06-1976
		NO 753984 A,B	03-06-1976
		PH 14474 A	07-08-1981
		SE 7513331 A	03-06-1976
		SE 8004364 A	11-06-1980
		SU 655311 A	30-03-1979
		SU 685146 A	05-09-1979
		US 4081547 A	28-03-1978
		US 4093742 A	06-06-1978
		US 4058612 A	15-11-1977
		US 4159335 A	26-06-1979
		ZA 7507457 A	24-11-1976
US 3281466 A	25-10-1966	DE 1543912 A	05-02-1970
		GB 1096369 A	
US 4251534 A	17-02-1981	BE 872956 A	21-06-1979
		CA 1099712 A	21-04-1981
		DE 2854727 A	28-06-1979
		DK 574978 A	23-06-1979
		FR 2412533 A	20-07-1979
		GB 2010821 A,B	04-07-1979
		IT 1102419 B	07-10-1985
		JP 54098780 A	03-08-1979
		LU 80720 A	22-01-1980
		NL 7812422 A	26-06-1979
US 4240979 A	23-12-1980	DE 2839462 A	27-03-1980
		AT 38 T	15-05-1981
		AU 522721 B	24-06-1982
		AU 5057679 A	20-03-1980
		DE 2960273 D	07-05-1981
		EP 0008768 A	19-03-1980
		HU 179118 B	28-08-1982

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/06611

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4240979 A		JP 1510053 C	26-07-1989
		JP 55038390 A	17-03-1980
		JP 63057418 B	11-11-1988
		US 4271171 A	02-06-1981
		ZA 7904787 A	24-09-1980
<hr/>			
EP 0193249 A	03-09-1986	AU 601145 B	06-09-1990
		AU 5410886 A	04-09-1986
		DK 88186 A	02-09-1986
		ES 552432 D	01-03-1988
		ES 8801896 A	16-05-1988
		GR 860542 A	25-06-1986
		JP 61218569 A	29-09-1986
		PH 23163 A	19-05-1989
		ZA 8601446 A	29-10-1986
<hr/>			